

**PCT**WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification<sup>6</sup> :C07D 307/20, 307/66, A61K 31/34,  
31/16, C07D 417/12, 405/12, 409/14,  
413/12, 409/12, 413/14, 407/12

A1

(11) International Publication Number:

WO 96/33184

(43) International Publication Date:

24 October 1996 (24.10.96)

(21) International Application Number:

PCT/US96/05475

(22) International Filing Date:

18 April 1996 (18.04.96)

(30) Priority Data:

08/424,819

19 April 1995 (19.04.95)

US

(71) Applicant: VERTEX PHARMACEUTICALS INCORPORATED [US/US]; 40 Allston Street, Cambridge, MA 02139-4211 (US).

(72) Inventor: TUNG, Roger, D.; 54 Richfield Road, Arlington, MA 02174 (US).

(74) Agents: HALEY, James, F., Jr. et al.; Fish &amp; Neave, 1251 Avenue of the Americas, New York, NY 10020 (US).

(81) Designated States: AL, AM, AT, AU, AZ, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).

**Published***With international search report.**Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.*

(54) Title: THF-CONTAINING SULFONAMIDE INHIBITORS OF ASPARTYL PROTEASE

## (57) Abstract

The present invention relates to a class of THF-containing sulfonamides which are aspartyl protease inhibitors. This invention also relates to pharmaceutical compositions comprising these compounds. The compounds and pharmaceutical compositions of this invention are particularly well suited for inhibiting HIV-1 and HIV-2 protease activity and consequently, may be advantageously used as anti-viral agents against the HIV-1 and HIV-2 viruses. This invention also relates to methods for inhibiting the activity of HIV aspartyl protease using the compounds of this invention.

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LJ	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

THF-CONTAINING SULFONAMIDE  
INHIBITORS OF ASPARTYL PROTEASE

TECHNICAL FIELD OF THE INVENTION

5     The present invention relates to a class of  
THF-containing sulfonamides which are aspartyl protease  
inhibitors. This invention also relates to  
pharmaceutical compositions comprising these compounds.  
The compounds and pharmaceutical compositions of this  
invention are particularly well suited for inhibiting  
10    HIV-1 and HIV-2 protease activity and consequently, may  
be advantageously used as anti-viral agents against the  
HIV-1 and HIV-2 viruses. This invention also relates  
to methods for inhibiting the activity of HIV aspartyl  
protease using the compounds of this invention.

15     BACKGROUND OF THE INVENTION

The human immunodeficiency virus ("HIV") is  
the causative agent for acquired immunodeficiency  
syndrome ("AIDS") -- a disease characterized by the  
destruction of the immune system, particularly of CD4<sup>+</sup>  
20    T-cells, with attendant susceptibility to opportunistic  
infections -- and its precursor AIDS-related complex  
("ARC") -- a syndrome characterized by symptoms such as  
persistent generalized lymphadenopathy, fever and  
weight loss.

25     As in the case of several other retroviruses,  
HIV encodes the production of a protease which carries

- 2 -

out post-translational cleavage of precursor polypeptides in a process necessary for the formation of infectious virions (S. Crawford et al., "A Deletion Mutation in the 5' Part of the pol Gene of Moloney Murine Leukemia Virus Blocks Proteolytic Processing of the gag and pol Polyproteins", J. Virol., 53, p. 899 (1985)). These gene products include pol, which encodes the virion RNA-dependent DNA polymerase (reverse transcriptase), an endonuclease, HIV protease, and gag, which encodes the core-proteins of the virion (H. Toh et al., "Close Structural Resemblance Between Putative Polymerase of a Drosophila Transposable Genetic Element 17.6 and pol gene product of Moloney Murine Leukemia Virus", EMBO J., 4, p. 1267 (1985); L.H. Pearl et al., "A Structural Model for the Retroviral Proteases", Nature, pp. 329-351 (1987); M.D. Power et al., "Nucleotide Sequence of SRV-1, a Type D Simian Acquired Immune Deficiency Syndrome Retrovirus", Science, 231, p. 1567 (1986)).

A number of synthetic anti-viral agents have been designed to target various stages in the replication cycle of HIV. These agents include compounds which block viral binding to CD4<sup>+</sup> T-lymphocytes (for example, soluble CD4), and compounds which interfere with viral replication by inhibiting viral reverse transcriptase (for example, didanosine and zidovudine (AZT)) and inhibit integration of viral DNA into cellular DNA (M.S. Hirsh and R.T. D'Aquila, "Therapy for Human Immunodeficiency Virus Infection", N.Eng.J.Med., 328, p. 1686 (1993)). However, such agents, which are directed primarily to early stages of viral replication, do not prevent the production of infectious virions in chronically infected cells. Furthermore, administration of some of these agents in effective amounts has led to cell-toxicity and unwanted

- 3 -

side effects, such as anemia and bone marrow suppression.

More recently, drug design efforts have been directed toward creating compounds which inhibit the formation of infectious virions by interfering with the processing of viral polyprotein precursors. Processing of these precursor proteins requires the action of virus-encoded proteases which are essential for replication (Kohl, N.E. et al. "Active HIV Protease is Required for Viral Infectivity" Proc. Natl. Acad. Sci. USA, 85, p. 4686 (1988)). The anti-viral potential of HIV protease inhibition has been demonstrated using peptidal inhibitors. Such peptidal compounds, however, are typically large and complex molecules that tend to exhibit poor bioavailability and are not generally consistent with oral administration. Accordingly, the need still exists for compounds that can effectively inhibit the action of viral proteases, for use as agents for preventing and treating chronic and acute viral infections. Such agents would be expected to act as effective therapeutic agents in their own right. In addition, since they act at a separate stage in the virus life cycle from previously described antiretroviral agents, the administration of a combination of agents would be expected to result in increased therapeutic efficacy.

#### SUMMARY OF THE INVENTION


The present invention provides a novel class of compounds, and pharmaceutically acceptable derivatives thereof, that are useful as inhibitors of aspartyl proteases, in particular, HIV aspartyl protease. These compounds can be used alone or in combination with other therapeutic or prophylactic agents, such as anti-virals, antibiotics,

- 4 -

immunomodulators or vaccines, for the treatment or prophylaxis of viral infection.

According to a preferred embodiment, the compounds of this invention are capable of inhibiting HIV viral replication in human CD<sub>4</sub><sup>+</sup> cells including T-cells, monocytic lines including macrophages and dendrocytes and other permissive cells. These compounds are useful as therapeutic and prophylactic agents to treat or prevent infection by HIV-1 and related viruses which may result in asymptomatic infection, AIDS-related complex ("ARC"), acquired immunodeficiency syndrome ("AIDS"), or similar disease of the immune system.

It is a principal object of this invention to provide a class of THF-containing sulfonamides which are aspartyl protease inhibitors, and particularly, HIV aspartyl protease inhibitors. This class of THF-containing sulfonamides is represented by formula I:



wherein:

each R<sup>1</sup> is independently selected from the group consisting of -C(O)-, -S(O)<sub>2</sub>-, -C(O)-C(O)-, -O-C(O)-, -O-S(O)<sub>2</sub>-, -NR<sup>2</sup>-S(O)<sub>2</sub>-, -NR<sup>2</sup>-C(O)- and -NR<sup>2</sup>-C(O)-C(O)-;

each Het is independently selected from the group consisting of C<sub>3</sub>-C<sub>7</sub> carbocycle; C<sub>6</sub>-C<sub>10</sub> aryl; phenyl fused with heterocycle; and heterocycle; wherein any member of said Het may be optionally substituted with one or more substituents selected from the group consisting of oxo, -OR<sup>2</sup>, -R<sup>2</sup>, -N(R<sup>2</sup>)(R<sup>2</sup>), -NHOH, -R<sup>2</sup>-OH, -CN, -CO<sub>2</sub>R<sup>2</sup>, -C(O)-N(R<sup>2</sup>)(R<sup>2</sup>), -S(O)<sub>2</sub>-N(R<sup>2</sup>)(R<sup>2</sup>), -N(R<sup>2</sup>)-

- 5 -

$C(O)-R^2$ ,  $-C(O)-R^2$ ,  $-S(O)_n-R^2$ ,  $-OCF_3$ ,  $-S(O)_n-R^6$ ,  $-N(R^2)-S(O)_2(R^2)$ , halo,  $-CF_3$ ,  $-NO_2$ ,  $-R^6$  and  $-O-R^6$ ;

5 each  $R^2$  is independently selected from the group consisting of H and  $C_1-C_3$  alkyl optionally substituted with  $R^6$ ;

10 each  $R^3$  is independently selected from the group consisting of H, Het,  $C_1-C_6$  alkyl and  $C_2-C_6$  alkenyl wherein any member of said  $R^3$ , except H, may be optionally substituted with one or more substituents selected from the group consisting of  $-OR^2$ ,  $-C(O)-NH-R^2$ ,  $-S(O)_n-N(R^2)(R^2)$ , Het,  $-CN$ ,  $-SR^2$ ,  $-CO_2R^2$ ,  $NR^2-C(O)-R^2$ ;

each  $n$  is independently 1 or 2;

15 each  $D$  and  $D'$  is independently selected from the group consisting of  $R^6$ ;  $C_1-C_3$  alkyl, which may be optionally substituted with one or more groups selected from  $-OR^2$ ,  $-R^3$ ,  $-O-R^6$ ,  $-S-R^6$  and  $R^6$ ;  $C_2-C_6$  alkenyl, which may be optionally substituted with one or more groups selected from the group consisting of  $-OR^2$ ,  $-R^3$ ,  $-O-R^6$  and  $R^6$ ; and  $C_3-C_6$  carbocycle, which may be optionally substituted with or fused with  $R^6$ ;

20 each  $E$  is independently selected from the group consisting of Het;  $-O-Het$ ;  $Het-Het$ ;  $-O-R^3$ ;  $-NR^2R^3$ ;  $C_1-C_6$  alkyl, which may be optionally substituted with one or more groups selected from the group consisting of  $R^4$  and Het; and  $C_2-C_6$  alkenyl, which may be optionally substituted with one or more groups selected from the group consisting of  $R^4$  and Het;

25 each  $R^4$  is independently selected from the group consisting of  $-OR^2$ ,  $-C(O)-NHR^2$ ,  $-S(O)_2-NHR^2$ , halo,  $-NR^2-C(O)-R^2$  and  $-CN$ ; and

30 each  $R^5$  is independently selected from the group consisting of H and  $C_1-C_6$  alkyl optionally substituted with aryl; and

35 each  $R^6$  is independently selected from the group consisting of aryl, carbocycle and heterocycle,

- 6 -

wherein said aryl, carbocycle or heterocycle may be optionally substituted with one or more groups selected from the group consisting of oxo, -OR<sup>s</sup>, -R<sup>s</sup>, -N(R<sup>s</sup>)(R<sup>s</sup>), -N(R<sup>s</sup>)-C(O)-R<sup>s</sup>, -R<sup>s</sup>-OH, -CN, -CO<sub>2</sub>R<sup>s</sup>, -C(O)-N(R<sup>s</sup>)(R<sup>s</sup>), halo and -CF<sub>3</sub>.

It is also an object of this invention to provide pharmaceutical compositions comprising the THF-containing sulfonamides of formula I and methods for their use as inhibitors of HIV aspartyl protease.

10

#### DETAILED DESCRIPTION OF THE INVENTION

In order that the invention herein described may be more fully understood, the following detailed description is set forth. In the description, the following abbreviations are used:

15	<u>Designation</u>	<u>Reagent or Fragment</u>
	Ac	acetyl
	Me	methyl
	Et	ethyl
	Bn	benzyl
20	Trityl	triphenylmethyl
	Asn	D- or L-asparagine
	Ile	D- or L-isoleucine
	Phe	D- or L-phenylalanine
	Val	D- or L-valine
25	Boc	tert-butoxycarbonyl
	Cbz	benzyloxycarbonyl (carbobenzyloxy)
	DCC	dicyclohexylcarbodiimide
	DBU	1,8-diazabicyclo(5.4.0)undec-7-ene
	DIC	diisopropylcarbodiimide
30	DIEA	diisopropylethylamine
	DMF	dimethylformamide
	DMSO	dimethylsulfoxide



- 7 -

	EDC	1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride
	EtOAc	ethyl acetate
	Fmoc	9-fluorenylmethoxycarbonyl
5	HOBt	1-hydroxybenzotriazole
	HOSu	1-hydroxysuccinimide
	iBu	iso-butyl
	NCA	N-carboxyanhydride
	t-Bu	tert-butyl
10	TFA	trifluoroacetic acid
	THP	tertrahdropyran
	THF	tetrahydrofuran
	TMSCl	chlorotrimethylsilane

15           The following terms are employed herein:  
          Unless expressly stated to the contrary, the terms "-SO<sub>2</sub>-" and "-S(O)<sub>2</sub>-" as used herein refer to a sulfone or sulfone derivative (i.e., both appended groups linked to the S), and not a sulfinat ester.

20           The term "backbone" refers to the structural representation of a compound of this invention, as set forth in the figures drawn in this application. The term "backbone" does not encompass the variables drawn in those figures.

25           For the compounds of formula I, and intermediates thereof, the stereochemistry of the explicitly shown hydroxyl is defined relative to D on the adjacent carbon atom, when the molecule is drawn in an extended zig-zag representation (such as that drawn for compounds of formula VI). If both OH and D reside  
30           on the same side of the plane defined by the extended backbone of the compound, the stereochemistry of the hydroxyl will be referred to as "syn". If OH and D reside on opposite sides of that plane, the

- 8 -

stereochemistry of the hydroxyl will be referred to as "anti".

As used herein, the term "alkyl", alone or in combination with any other term, refers to a  
5 straight-chain or branch-chain saturated aliphatic hydrocarbon radical containing the specified number of carbon atoms, or where no number is specified, preferably from 1-10 and more preferably from 1-5 carbon atoms. Examples of alkyl radicals include, but  
10 are not limited to, methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, sec-butyl, tert-butyl, pentyl, isoamyl, n-hexyl and the like.

The term "alkenyl", alone or in combination with any other term, refers to a straight-chain or  
15 branched-chain mono- or poly-unsaturated aliphatic hydrocarbon radical containing the specified number of carbon atoms, or where no number is specified, preferably from 2-10 carbon atoms and more preferably, from 2-6 carbon atoms. Examples of alkenyl radicals  
20 include, but are not limited to, ethenyl, E- and Z-propenyl, isopropenyl, E- and Z-butenyl, E- and Z-isobutenyl, E- and Z-pentenyl, E- and Z-hexenyl, E,E-, E,Z-, Z,E- and Z,Z-hexadienyl and the like.

The term "aryl", alone or in combination with  
25 any other term, refers to a carbocyclic aromatic radical (such as phenyl or naphthyl) containing the specified number of carbon atoms, preferably from 6-14 carbon atoms, and more preferably from 6-10 carbon atoms. Examples of aryl radicals include, but are not  
30 limited to phenyl, naphthyl, indenyl, indanyl, azulenyl, fluorenyl, anthracenyl and the like.

The term "cycloalkyl", alone or in combination with any other term, refers to a cyclic saturated hydrocarbon radical containing the specified  
35 number of carbon atoms, preferably from 3-7 carbon

- 9 -

atoms. Examples of cycloalkyl radicals include, but are not limited to, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl and the like.

5 The term "cycloalkenyl", alone or in combination with any other term, refers to a cyclic hydrocarbon radical containing the specified number of carbon atoms with at least one endocyclic carbon-carbon bond. Where no number of carbon atoms is specified, a cycloalkenyl radical preferably has from 5-7 carbon  
10 atoms. Examples of cycloalkenyl radicals include, but are not limited to, cyclopentenyl, cyclohexenyl, cyclopentadienyl and the like.

The term "THF" refers to a tetrahydrofuran ring attached at any ring carbon resulting in a stable  
15 structure, but preferably attached at the 3-position of the tetrahydrofuran ring (i.e., tetrahydrofuran-3-yl). Preferably, the chiral carbon of THF is in the (S) configuration.

The term "carbocycle" refers to a stable  
20 non-aromatic 3- to 8-membered carbon ring radical which may be saturated, mono-unsaturated or poly-unsaturated. The carbocycle may be attached at any endocyclic carbon atom which results in a stable structure. Preferred carbocycles have 5-6 carbons. Examples of carbocycle  
25 radicals include, but are not limited to, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, cycloheptyl, cyclopentenyl, cyclohexenyl, cyclopentadienyl and the like.

The term "heterocycle", unless otherwise  
30 defined herein, refers to a stable 3-7 membered monocyclic heterocyclic ring or 8-11 membered bicyclic heterocyclic ring which is either saturated or unsaturated, and which may be optionally benzofused if monocyclic. Each heterocycle consists of one or more  
35 carbon atoms and from one to four heteroatoms selected

- 10 -

from the group consisting of nitrogen, oxygen and sulfur. As used herein, the terms "nitrogen and sulfur heteroatoms" include any oxidized form of nitrogen and sulfur, and the quaternized form of any basic nitrogen.

5 In addition, any ring nitrogen may be optionally substituted with a substituent  $R^2$ , as defined herein for compounds of formula I. A heterocycle may be attached at any endocyclic carbon or heteroatom which results in the creation of a stable structure. A heterocycle may

10 be attached at any endocyclic carbon or heteroatom which results in the creation of a stable structure. Preferred heterocycles include 5-7 membered monocyclic heterocycles and 8-10 membered bicyclic heterocycles. Preferred heterocycles defined above include, for

15 example, benzimidazolyl, imidazolyl, imidazolinoyl, imidazolidinyl, quinolyl, isoquinolyl, indolyl, indazolyl, indazolinolyl, perhydropyridazyl, pyridazyl, pyridyl, pyrrolyl, pyrrolinyl, pyrrolidinyl, pyrazolyl, pyrazinyl, quinoxolyl, piperidinyl, pyranyl,

20 pyrazolinyl, piperazinyl, pyrimidinyl, pyridazinyl, morpholinyl, thiamorpholinyl, furyl, thienyl, triazolyl, thiazolyl,  $\beta$ -carboline, tetrazolyl, thiazolidinyl, benzofuranoyl, thiamorpholinyl sulfone, oxazolyl, benzoxazolyl, oxopiperidinyl, oxopyrrolidinyl,

25 oxoazepinyl, azepinyl, isoxazolyl, isothiazolyl, furazanyl, tetrahydropyranyl, tetrahydrofuranyl, thiazolyl, thiadiazoyl, dioxolyl, dioxinyl, oxathieryl, benzodioxolyl, dithieryl, thiophenyl, tetrahydrothiophenyl, dioxanyl, dioxolanyl,

30 tetrahydrofurotetrahydrofuranyl, tetrahydropyranotetrahydrofuranyl, tetrahydrofurodihydrofuranyl, , tetrahydropyranodihydrofuranyl, dihydropyranyl, dihydrofuranyl, dihydrofurotetrahydrofuranyl,

- 11 -

dihydropyranotetrahydrofuranyl, sulfolanyl and the like.

The term "halo" refers to a radical of fluorine, chlorine, bromine or iodine.

5           The terms "HIV protease" and "HIV aspartyl protease" are used interchangeably and refer to the aspartyl protease encoded by the human immunodeficiency virus type 1 or 2. In a preferred embodiment of this invention, these terms refer to the human  
10 immunodeficiency virus type 1 aspartyl protease.

          The term "anti-viral agent" or "anti-retroviral agent" refers to a compound or drug which possesses viral inhibitory activity. Such agents include reverse transcriptase inhibitors (including  
15 nucleoside and non-nucleoside analogs) and protease inhibitors. Preferably the protease inhibitor is an HIV protease inhibitor. Examples of nucleoside analog reverse transcriptase inhibitors include, but are not limited to, zidovudine (AZT), dideoxycytidine (ddC),  
20 didanosine (ddI), stavudine (d4T), 3TC, 935U83, 1592U89 and 524W91. Examples of non-nucleoside analog reverse transcriptase inhibitors include, but are not limited to delavirdine (U90) and nevirapine. Examples of HIV  
25 protease inhibitors include, but are not limited to, saquinavir (Ro 31-8959), MK 639, ABT 538 (A80538), AG 1343, XM 412, XM 450, BMS 186318 and CPG 53,437.

          The term "leaving group" or "LG" refers to groups readily displaceable by a nucleophile, such as an amine, alcohol, phosphorous or thiol nucleophile or  
30 their respective anions. Such leaving groups are well known and include carboxylates, N-hydroxysuccinimide, N-hydroxybenzotriazole, halogen (halides), triflates, tosylates, mesylates, alkoxy, thioalkoxy, phosphinates, phosphonates and the like. Other potential  
35 nucleophiles include organometallic reagents known to

- 12 -

those skilled in the art. In addition, the term "leaving group" or "LG" is meant to encompass leaving group precursors (i.e., moieties that can be easily converted to a leaving group upon simple synthetic procedures such as alkylation, oxidation or protonation). Such leaving group precursors and methods for converting them to leaving groups are well known to those of ordinary skill in the art. Leaving group precursors include, for instance, secondary and tertiary amines. By way of example, the moiety -N(R<sub>3</sub>)(R<sub>4</sub>), while not itself a leaving group, is encompassed by the term "leaving group" or "LG" because it can be readily converted to a leaving group such as -N<sup>+</sup>CH<sub>3</sub>(R<sub>3</sub>)(R<sub>4</sub>).

The term "protecting group" refers to a suitable chemical group which may be attached to a functional group and removed at a later stage to reveal the intact functional group. Examples of suitable protecting groups for various functional groups are described in T.W. Greene and P.G.M. Wuts, Protective Groups in Organic Synthesis, 2d. Ed., John Wiley and Sons (1991); L. Fieser and M. Fieser, Fieser and Fieser's Reagents for Organic Synthesis, John Wiley and Sons (1994); L. Paquette, ed. Encyclopedia of Reagents for Organic Synthesis, John Wiley and Sons (1995).

The term "silyl" refers to a trisubstituted silicon radical in which the substituents are independently C<sub>1</sub>-C<sub>4</sub> alkyl, C<sub>6</sub>-C<sub>7</sub> aryl or C<sub>3</sub>-C<sub>7</sub> carbocycle. Examples of silyl groups include, but are not limited to, trimethylsilyl, triethylsilyl, triisopropylsilyl, t-butyl dimethylsilyl, t-butyl diisopropylsilyl, t-butyl diphenylsilyl, triphenylsilyl, cyclohexyldimethylsilyl and the like.

The term "pharmaceutically effective amount" refers to an amount effective in treating HIV infection

- 13 -

in a patient either as monotherapy or in combination with other agents. The term "treating" as used herein refers to the alleviation of symptoms of a particular disorder in a patient or the improvement of an

5 ascertainable measurement associated with a particular disorder. Specifically, with respect to HIV, effective treatment using the compounds and compositions of this invention would result in an improvement in an HIV associated ascertainable measurement. Such

10 measurements include, but are not limited to, reduction in viral load in plasma or another defined tissue compartment as measured by, e.g. RT-PCR or branched-chain DNA PCR or culturable virus measurements,  $\beta$ -2 microglobulin or p24 levels, number of CD<sub>4</sub><sup>+</sup> cells or

15 ratio of CD<sub>4</sub><sup>+</sup>/CD<sub>8</sub><sup>+</sup> cells, or functional markers such as improvement in quality of life, ability to carry out normal functions, reduction of dementia or immunosuppression-related effects including, but not

20 limited to, opportunistic infections and tumors. The term "prophylactically effective amount" refers to an amount effective in preventing HIV infection in a patient. As used herein, the term "patient" refers to a mammal, including a human.

The term "pharmaceutically acceptable carrier or adjuvant" refers to a carrier or adjuvant that may

25 be administered to a patient, together with a compound of this invention, and which does not destroy the pharmacological activity thereof and is nontoxic when administered in doses sufficient to deliver a

30 therapeutic amount of the antiretroviral agent.

The term "point of attachment" refers to the atom through which a moiety is attached to a specified structure.

The term "substituted", whether express or

35 implied and whether preceded by the term "optionally"

- 14 -

or not, refers to the replacement of one or more hydrogen radicals in a given structure with the radical of a specified substituent. When more than one position in a given structure may be substituted with a substituent selected from a specified group, the substituents may be either the same or different at every position. Typically, when a structure may be optionally substituted, 0-3 substitutions are preferred, and 0-1 substitution is most preferred. Most preferred substituents are those which enhance protease inhibitory activity or intracellular antiviral activity in permissive mammalian cells or immortalized mammalian cell lines, or which enhance deliverability by enhancing solubility characteristics or enhancing pharmacokinetic or pharmacodynamic profiles as compared to the unsubstituted compound. Other most preferred substituents include those used in the compounds shown in Table I.

As used herein, the compounds of this invention, including the compounds of formula I, are defined to include pharmaceutically acceptable derivatives or prodrugs thereof. A "pharmaceutically acceptable derivative or prodrug" means any pharmaceutically acceptable salt, ester, salt of an ester, or other derivative of a compound of this invention which, upon administration to a recipient, is capable of providing (directly or indirectly) a compound of this invention or an inhibitorily active metabolite or residue thereof. Particularly favored derivatives and prodrugs are those that increase the bioavailability of the compounds of this invention when such compounds are administered to a mammal (e.g., by allowing an orally administered compound to be more readily absorbed into the blood) or which enhance delivery of the parent compound to a biological



- 15 -

compartment (e.g., the brain or lymphatic system) relative to the parent species. Preferred prodrugs include derivatives where a group which enhances aqueous solubility or active transport through the gut membrane is appended to the explicitly shown hydroxyl in formula (I) or to "E" in formula (I).

Pharmaceutically acceptable salts of the compounds of this invention include those derived from pharmaceutically acceptable inorganic and organic acids and bases. Examples of suitable acids include hydrochloric, hydrobromic, sulfuric, nitric, perchloric, fumaric, maleic, phosphoric, glycollic, lactic, salicylic, succinic, p-toluenesulfonic, tartaric, acetic, citric, methanesulfonic, ethanesulfonic, formic, benzoic, malonic, naphthalene-2-sulfonic and benzenesulfonic acids. Preferred acids include hydrochloric, sulfuric, methanesulfonic and ethanesulfonic acids. Methanesulfonic acid is most preferred. Other acids, such as oxalic, while not in themselves pharmaceutically acceptable, may be employed in the preparation of salts useful as intermediates in obtaining the compounds of the invention and their pharmaceutically acceptable acid addition salts.

Salts derived from appropriate bases include alkali metal (e.g., sodium), alkaline earth metal (e.g., magnesium), ammonium and N-(C<sub>1-4</sub> alkyl)<sub>4</sub><sup>+</sup> salts.

The term "thiocarbamates" refers to compounds containing the functional group N-SO<sub>2</sub>-O.

The compounds of this invention contain one or more asymmetric carbon atoms and thus occur as racemates and racemic mixtures, single enantiomers, diastereomeric mixtures and individual diastereomers. All such isomeric forms of these compounds are expressly included in the present invention. Each stereogenic carbon may be of the R or S configuration.

- 16 -

The explicitly shown hydroxyl is also preferred to be syn to D, in the extended zig-zag conformation between the nitrogens shown in compounds of formula I.

Combinations of substituents and variables  
5 envisioned by this invention are only those that result in the formation of stable compounds. The term "stable", as used herein, refers to compounds which possess stability sufficient to allow manufacture and which maintains the integrity of the compound for a  
10 sufficient period of time to be useful for the purposes detailed herein (e.g., therapeutic or prophylactic administration to a mammal or for use in affinity chromatography applications). Typically, such compounds are stable at a temperature of 40°C or less,  
15 in the absence of moisture or other chemically reactive conditions, for at least a week.


The compounds of the present invention may be used in the form of salts derived from inorganic or organic acids. Included among such acid salts, for  
20 example, are the following: acetate, adipate, alginate, aspartate, benzoate, benzenesulfonate, bisulfate, butyrate, citrate, camphorate, camphorsulfonate, cyclopentanepropionate, digluconate, dodecylsulfate, ethanesulfonate, fumarate, glucohepta-  
25 noate, glycerophosphate, hemisulfate, heptanoate, hexanoate, hydrochloride, hydrobromide, hydroiodide, 2-hydroxyethanesulfonate, lactate, maleate, methanesulfonate, 2-naphthalenesulfonate, nicotinate, oxalate, palmoaté, pectinate, persulfate, 3-  
30 phenylpropionate, picrate, pivalate, propionate, succinate, tartrate, thiocyanate, tosylate and undecanoate.

This invention also envisions the quaternization of any basic nitrogen-containing groups  
35 of the compounds disclosed herein. The basic nitrogen

- 17 -

can be quaternized with any agents known to those of ordinary skill in the art including, for example, lower alkyl halides, such as methyl, ethyl, propyl and butyl chlorides, bromides and iodides; dialkyl sulfates including dimethyl, diethyl, dibutyl and diamyl sulfates; long chain halides such as decyl, lauryl, myristyl and stearyl chlorides, bromides and iodides; and aralkyl halides including benzyl and phenethyl bromides. Water or oil-soluble or dispersible products may be obtained by such quaternization.

The THF-containing sulfonamides of this invention are those of formula I:



wherein:

each  $\text{R}^1$  is independently selected from the group consisting of  $-\text{C}(\text{O})-$ ,  $-\text{S}(\text{O})_2-$ ,  $-\text{C}(\text{O})-\text{C}(\text{O})-$ ,  $-\text{O}-\text{C}(\text{O})-$ ,  $-\text{O}-\text{S}(\text{O})_2-$ ,  $-\text{NR}^2-\text{S}(\text{O})_2-$ ,  $-\text{NR}^2-\text{C}(\text{O})-$  and  $-\text{NR}^2-\text{C}(\text{O})-\text{C}(\text{O})-$ ; preferably each  $\text{R}^1$  is  $-\text{O}-\text{C}(\text{O})-$  or  $-\text{C}(\text{O})-$ ; more preferably each  $\text{R}^1$  is  $-\text{O}-\text{C}(\text{O})-$ ;

each Het is independently selected from the group consisting of  $\text{C}_3\text{-C}_7$  carbocycle;  $\text{C}_6\text{-C}_{10}$  aryl; phenyl fused with heterocycle; and heterocycle; wherein any member of said Het may be optionally substituted with one or more substituents selected from the group consisting of oxo,  $-\text{OR}^2$ ,  $-\text{R}^2$ ,  $-\text{N}(\text{R}^2)(\text{R}^2)$ ,  $-\text{NHOH}$ ,  $-\text{R}^2\text{-OH}$ ,  $-\text{CN}$ ,  $-\text{CO}_2\text{R}^2$ ,  $-\text{C}(\text{O})-\text{N}(\text{R}^2)(\text{R}^2)$ ,  $-\text{S}(\text{O})_2-\text{N}(\text{R}^2)(\text{R}^2)$ ,  $-\text{N}(\text{R}^2)-\text{C}(\text{O})-\text{R}^2$ ,  $-\text{C}(\text{O})-\text{R}^2$ ,  $-\text{S}(\text{O})_n-\text{R}^2$ ,  $-\text{OCF}_3$ ,  $-\text{S}(\text{O})_n-\text{R}^6$ ,  $-\text{N}(\text{R}^2)-\text{S}(\text{O})_2(\text{R}^2)$ , halo,  $-\text{CF}_3$ ,  $-\text{NO}_2$ ,  $-\text{R}^6$  and  $-\text{O}-\text{R}^6$ ;

each  $\text{R}^2$  is independently selected from the group consisting of H and  $\text{C}_1\text{-C}_3$  alkyl optionally substituted with  $\text{R}^6$ ;

- 18 -

each  $R^3$  is independently selected from the group consisting of H, Het,  $C_1-C_6$  alkyl and  $C_2-C_6$  alkenyl wherein any member of said  $R^3$ , except H, may be optionally substituted with one or more substituents selected from the group consisting of  $-OR^2$ ,  $-C(O)-NH-R^2$ ,  
5  $-S(O)_n-N(R^2)(R^2)$ , Het,  $-CN$ ,  $-SR^2$ ,  $-CO_2R^2$ ,  $NR^2-C(O)-R^2$ ;

each  $n$  is independently 1 or 2;

each  $D$  and  $D'$  is independently selected from the group consisting of  $R^6$ ;  $C_1-C_6$  alkyl, which may be optionally substituted with one or more groups selected  
10 from  $-OR^2$ ,  $-R^3$ ,  $-O-R^6$  and  $R^6$ ;  $C_2-C_4$  alkenyl, which may be optionally substituted with one or more groups selected from the group consisting of  $-OR^2$ ,  $-R^3$ ,  $-O-R^6$  and  $R^6$ ;  $C_3-C_6$  carbocycle, which may be optionally substituted with  
15 or fused with  $R^6$ ; and  $C_3-C_6$  cycloalkenyl, which may be optionally substituted with or fused with  $R^6$ ; preferably each  $D$  is independently  $C_1-C_3$  alkyl, which may be optionally substituted with one or more Het; more preferably  $D$  is  $C_1-C_3$  alkyl, which may be optionally  
20 substituted with one group selected from  $C_6-C_{10}$  aryl and  $C_3-C_6$  cycloalkyl; even more preferably  $D$  is selected from benzyl, isobutyl, cyclopentylmethyl, and cyclohexylmethyl; and most preferably,  $D$  is benzyl or isobutyl; preferably each  $D'$  is independently selected  
25 from the group consisting of  $C_1-C_6$  alkyl optionally substituted with  $R^6$ ; and more preferably  $D'$  is selected from the group consisting of  $C_1-C_4$  alkyl optionally substituted with one 3-6 membered carbocycle or one 5-6 membered heterocycle; and most preferably,  $D'$  is  
30 selected from the group consisting of isobutyl, cyclopentylmethyl and cyclohexylmethyl;

each  $E$  is independently selected from the group consisting of Het;  $-O-Het$ ;  $Het-Het$ ;  $-O-R^3$ ;  $-NR^3R^3$ ;  $C_1-C_6$  alkyl, which may be optionally substituted with  
35 one or more groups selected from the group consisting

- 19 -

of R<sup>4</sup> and Het; C<sub>2</sub>-C<sub>6</sub> alkenyl, which may be optionally substituted with one or more groups selected from the group consisting of R<sup>4</sup> and Het; and phenyl fused with 5-6 membered heterocycle; preferably each E is Het and more preferably, E is phenyl substituted with one or more substituents selected from the group consisting of -OH, -OCH<sub>3</sub>, -NH<sub>2</sub>, -NHCOCH<sub>3</sub>, -SH, and -CH<sub>3</sub>; or phenyl fused with 5-6 membered heterocycle, and most preferably, E is phenyl substituted with -NH<sub>2</sub> (preferably in the meta- or para-position);

each R<sup>4</sup> is independently selected from the group consisting of -OR<sup>2</sup>, -C(O)-NHR<sup>2</sup>, -S(O)<sub>2</sub>-NHR<sup>2</sup>, halo, -NR<sup>2</sup>-C(O)-R<sup>2</sup> and -CN;

each R<sup>5</sup> is independently selected from the group consisting of H and C<sub>1</sub>-C<sub>4</sub> alkyl optionally substituted with aryl; preferably each R<sup>5</sup> is independently selected from the group consisting of H and C<sub>1</sub>-C<sub>3</sub> alkyl; and

each R<sup>6</sup> is independently selected from the group consisting of aryl, carbocycle and heterocycle, wherein said carbocycle or heterocycle may be optionally substituted with one or more groups selected from the group consisting of oxo, -OR<sup>5</sup>, -R<sup>5</sup>, -N(R<sup>5</sup>)(R<sup>5</sup>), -N(R<sup>5</sup>)-C(O)-R<sup>5</sup>, -R<sup>5</sup>-OH, -CN, -CO<sub>2</sub>R<sup>5</sup>, -C(O)-N(R<sup>5</sup>)(R<sup>5</sup>), halo and -CF<sub>3</sub>; preferably each R<sup>6</sup> is independently selected from the group consisting of 3-6 membered carbocycle and 5-6 membered heterocycle, wherein said heterocycle or carbocycle may be optionally substituted with one or more groups selected from the group consisting of oxo, -OR<sup>5</sup>, -R<sup>5</sup>, -N(R<sup>5</sup>)(R<sup>5</sup>), -N(R<sup>5</sup>)-C(O)-R<sup>5</sup>, -R<sup>5</sup>-OH, -CN, -CO<sub>2</sub>R<sup>5</sup>, -C(O)-N(R<sup>5</sup>)(R<sup>5</sup>), halo and -CF<sub>3</sub>.

Except where expressly noted to the contrary, the term "[variable]" as defined for formula I" refers to the definitions shown directly above.

- 20 -


Preferred compounds of formula I include those compounds having at least one variable defined as the preferred, more preferred, even more preferred or most preferred definition above. More preferred

5 compounds of formula I include those compounds having at least two to three variables defined independently as the preferred, more preferred, even more preferred or most preferred definitions above. Most preferred



















10 compounds of formula I include those compounds having at least four to five variables independently defined as the preferred, more preferred, even more preferred or most preferred definitions above.

Table I illustrates preferred compounds of this invention:



















- 21 -






































#	THF	OH	D	D'	E
35	S	syn			
37	S	syn			
48	S	syn			
51	S	syn			



















52	R,S	syn			
53	S	syn			
60	S	syn			
66	S	syn			
69	S	syn			
86	R,S	syn			







































88	S	syn			
91	S	syn			
93	R	syn			
94	R	syn			
95	S	syn			
99	S	syn			



















100	S	syn			
101	S	syn		CH <sub>3</sub>	
112	S	syn			
113	S	syn			
116	S	syn			
123	S	syn			



















124	S	syn			
125	S	syn			
132	S	syn			
133	S	syn			
134	S	syn			
135	S	syn			

136	S	syn			
137	S	syn			
138	S	syn			
140	S	syn			
144	S	syn			
145	S	syn			



















148	S	syn			
149	S	syn			
150	S	syn			
151	S	syn			
152	S	syn			
157	S	syn			



















158	S	syn			
159	S	syn			
160	S	syn			
161	S	syn			
165	S	syn			
167	S	syn			



















168	S	syn			
169	S	syn			
170	S	syn			
171	S	syn			
172	S	syn			
173	S	syn			



















174	S	syn			
175	S	syn			
176	S	syn			
180	S	syn			
181	S	syn			
182	S	syn			







































183	S	syn			
195	S	syn			
196	S	syn			
197	S	syn			
198	S	syn			
199	S	syn			



















200	S	syn			
201	S	syn			
202	S	syn			
203	S	anti			
204	S	syn			
205	S	syn			




206	S	syn			
207	S	anti			
208	S	syn			
209	S	syn			
210	S	syn			
211	S	syn			

212	S	syn			
213	S	syn			
214	S	syn			
215	S	syn			
216	S	syn			
217	S	syn			

218	S	syn			
219	S	syn			
220	S	syn			
221	S	syn			
222	S	syn			
223	S	syn			

224	S	syn			
225	S	syn			
226	S	syn			
227	S	syn			
228	S	syn			
229	S	syn			

230	S	syn			
231	S	syn			
232	S	syn			
233	S	syn			
234	S	syn			
235	S	syn			

236	S	syn			
-----	---	-----	---	--	---



- 39 -

More preferred compounds of this invention are selected from the group consisting of compound 35; compound 37; compound 48; compounds 52; compound 60; compound 66; compounds 86; compound 88; compound 91; compound 93; compound 94; compound 95; compound 99; compound 100; compound 112; compound 113; compound 116; compound 124; compound 125; compound 132; compound 134; compound 135; compound 138; compound 140; compound 144; compound 145; compound 148; compound 149; compound 150; compound 151; compound 152; compound 157; compound 158; compound 159; compound 160; compound 165; compound 167; compound 168; compound 169; compound 170; compound 171; compound 173; compound 175; compound 176; compound 180; compound 181; compound 182; compound 183; compound 195; compound 196; compound 197; compound 198; compound 200; compound 201; compound 202; compound 203; compound 204; compound 205; compound 206; compound 208; compound 209; compound 210; compound 211; compound 212; compound 213; compound 216; compound 217; compound 218; compound 219; compound 220; compound 221; compound 222; compound 224; compound 227; and compound 233, wherein each compound has the formula shown in Table I.

Even more preferred compounds of this invention are selected from the group consisting of: compound 48; compound 100; compound 116, compound 140; compound 148; compound 158; compound 160; compound 168; compound 169; compound 171; compound 173; compound 175; compound 176; compound 180; compound 181; compound 195; compound 197; compound 198; compound 202; compound 206; compound 211; compound 216; compound 217; compound 219 and compound 220, wherein each compound has the formula shown in Table I.

The most preferred compounds of this invention are selected from the group consisting of:

- 40 -


compound 140; compound 168; compound 169; compound 171; compound 175; compound 216; and compound 217, wherein each compound has the formula shown in Table I.

5 The THF-containing sulfonamides of this invention may be synthesized using conventional techniques. Advantageously, these compounds may be conveniently synthesized from readily available starting materials.

10 The compounds of this invention are among the most readily synthesized HIV protease inhibitors known. Previously described HIV protease inhibitors often contain four or more chiral centers, numerous peptide linkages and/or require air-sensitive reagents (such as organometallic complexes) to effect their synthesis.

15 The relative ease with which the compounds of this invention can be synthesized represents an enormous advantage in the large scale production of these compounds.

20 In general, the THF-containing sulfonamides of this invention are conveniently obtained from  $\alpha$ -amino acids and derivatives thereof having the general formula II:



25 wherein W is hydrogen or P; P is a suitable amino protecting group; Q is hydrogen, benzyl or A-R<sup>1</sup>-; Y is -C(O)OH, -C(O)H, or -CH<sub>2</sub>OH; and D and A-R<sup>1</sup>- are as defined above for the compounds of formula I. W and Q may also be taken together with the nitrogen to which they are attached to form a heterocycle. An example of

30 such a construction is phthalimide. Suitable amino protecting groups are described in numerous references, including T.W. Greene and P.G.M. Wuts, Protective

- 41 -

Groups in Organic Synthesis, 2d Ed., John Wiley and Sons (1991); L. Fieser and M. Fieser, Fieser and Fieser's Reagents for Organic Synthesis, John Wiley and Sons (1994); and L. Paquette, ed. Encyclopedia of Reagents for Organic Synthesis, John Wiley and Sons (1995). Examples of such amino protecting groups include, but are not limited to, Boc, Cbz and Alloc. Alternatively, the amine may be protected as an alkyl derivative such as N,N-dibenzyl or trityl. Such  $\alpha$ -amino acid derivatives are often commercially available or may be conveniently prepared from commercially available  $\alpha$ -amino acid derivatives using known techniques. Although this invention envisions the use of racemic mixtures of such starting materials, a single enantiomer (preferably in the S configuration) is preferred.

Using known techniques, the  $\alpha$ -amino acid derivative of general formula  $P-N(Q)-CH(D)-COOH$  may be readily converted to an amino ketone derivative of general formula  $P-N(Q)-CH(D)-CO-CH_2-X$ , wherein P, Q and D are as defined for compounds of formula II and X is a leaving group which suitably activates the  $\alpha$ -carbon (i.e., increases the susceptibility of the methylene to nucleophilic attack). Suitable leaving groups are well known in the art and include halides, dialkyl sulfonium salts and sulfonates, such as methanesulfonate, trifluoromethanesulfonate or 4-toluenesulfonate. X may also be a hydroxyl which is converted in situ to a leaving group (e.g., by treatment with a trialkyl- or triarylphosphine in the presence of a dialkylazodicarboxylate). Methods for the formation of such amino ketone derivatives also are well known to those of skill in the art (see, for example, S.J. Fittkau, J. Prakt. Chem., 315, p. 1037 (1973)). In addition, certain amino ketone derivatives are

- 42 -

commercially available (e.g., from Bachem Biosciences, Inc., Philadelphia, Pennsylvania).

The amino ketone derivative may then be reduced to the corresponding amino alcohol, represented by the formula  $P-N(Q)-CH(D)-CH(OH)-CH_2-X$ , wherein P, Q and D are as defined above for compounds of formula II and X is a leaving group as defined above.


Alternatively, the amino ketone derivative can be reduced to the corresponding alcohol later in the synthetic scheme. Many techniques for reduction of amino ketone derivatives such as  $P-N(Q)-CH(D)-CO-CH_2-X$  are well known to those of ordinary skill in the art (G.J. Quallich and T.M. Woodall, Tetrahedron Lett., 34, p. 785 (1993) and references cited therein; and Larock, R.C. "Comprehensive Organic Transformations", pp. 527-547, VCH Publishers, Inc.® 1989 and references cited therein). A preferred reducing agent is sodium borohydride. The reduction reaction is typically conducted at a temperature of from about  $-40^{\circ}\text{C}$  to about  $40^{\circ}\text{C}$  (preferably, at about  $0^{\circ}\text{C}$  to about  $20^{\circ}\text{C}$ ), in a suitable solvent system such as, for example, aqueous or neat tetrahydrofuran or a lower alcohol, such as methanol or ethanol. Although this invention envisions both stereospecific and non-stereospecific reduction of the amino ketone derivative  $P-N(Q)-CH(D)-CO-CH_2-X$ , stereoselective reduction is preferred. Stereoselective reduction may be accomplished by use of chiral reagents known in the art or by the use of an achiral reducing agent on a chiral substrate. In the present invention, stereoselective reduction may be conveniently achieved, for instance, under non-chelating reducing conditions, where chiral induction of the newly formed hydroxyl group is set by the stereochemistry of the D group (i.e., Felkin-Ahn addition of hydride). We particularly prefer stereoselective reductions wherein

- 43 -

the resulting hydroxyl is syn to D. We have found that when the hydroxyl group is syn to D, the final sulfonamide product is a more potent HIV protease inhibitor than its anti diastereomer.

5 The hydroxyl group of the amino alcohol may optionally be protected by any known oxygen protecting group (such as trialkylsilyl, benzyl, acetal or alkyloxymethyl) to yield a protected amino alcohol having the formula  $P-N(Q)-CH(D)-C(OR')-CH_2-X$ , wherein P,  
10 Q and D are as defined for compounds of formula II, X is a leaving group as defined above and  $R'$  is H or a suitable hydroxy protecting group. Several useful protecting groups are described in T.W. Greene and P.G.M. Wuts, Protective Groups in Organic Synthesis, 2d  
15 Ed., John Wiley and Sons (1991); L. Fieser and M. Fieser, Fieser and Fieser's Reagents for Organic Synthesis, John Wiley and Sons (1994); and L. Paquette, ed. Encyclopedia of Reagents for Organic Synthesis, John Wiley and Sons (1995).

20           The amino alcohol may then be reacted with a nucleophilic amine compound to form an intermediate of formula III:




wherein W, Q and D are as defined in formula II, R' is H or any suitable oxygen protecting group and L is either D' (as described for compounds of formula I) or hydrogen.

Alternatively, an amino acid derivative may be reacted with a nucleophilic nitro compound (e.g., a nitromethane anion or a derivative thereof) which can

- 44 -

be reduced in one or more steps to yield an intermediate of formula III.

In a particularly advantageous synthetic scheme, simultaneous activation of the methylene and protection of the alcohol may be accomplished by forming an N-protected amino epoxide from the oxygen and its adjacent methylene to give an intermediate of formula IV:



wherein W, Q and D are as defined above for compounds of formula II. Suitable solvent systems for preparing the N-protected amino epoxide include ethanol, methanol, isopropanol, tetrahydrofuran, dioxane, dimethylformamide and the like (including mixtures thereof). Suitable bases for producing the epoxide include alkali metal hydroxides, potassium t-butoxide, DBU and the like. A preferred base is potassium hydroxide.

Alternatively, the N-protected amino epoxide may be prepared by reacting an (alkylthio) or (phenylthio)acetic acid dianion with a cyclic N-carboxyanhydride of a protected  $\alpha$ -amino acid (such as BOC-Phe-NCA, available from Propeptide). A preferred acetic acid dianion is (methylthio)acetic acid dianion. The resulting amino ketone may then be reduced (e.g., with sodium borohydride). The resulting amino alcohol is readily converted to the amino epoxide by quaternization (e.g., with methyl iodide) followed by ring closure (using, for example, sodium hydride).

Reaction of the N-protected amino epoxide (or other suitably activated intermediate) with an amine is carried out neat, i.e. in the absence of solvent, or in the presence of a polar solvent such as lower alkanols,

- 45 -

water, dimethylformamide or dimethylsulfoxide. The reaction can be carried out conveniently between about -30°C and 120°C, preferably between about -5°C and 100°C. Alternatively, the reaction may be carried out in the presence of an activating agent, such as activated alumina in an inert solvent, preferably an ether, such as diethyl ether, tetrahydrofuran, dioxane, or tert-butyl methyl ether, conveniently from about room temperature to about 110°C, as described by Posner and Rogers, J. Am. Chem. Soc., 99, p. 8208 (1977). Other activating reagents include lower trialkylaluminum species, such as triethylaluminum, or dialkylaluminum halide species, such as diethylaluminum chloride (Overman and Flippin, Tetrahedron Letters, p. 195 (1981)). Reactions involving these species are conveniently carried out in inert solvents such as dichloromethane, 1,2-dichloroethane, toluene, or acetonitrile between about 0°C and about 110°C. Further methods of displacing leaving groups, or opening epoxides with amines or their equivalents such as azides or trimethylsilyl cyanide (Gassman and Guggenheim, J. Am. Chem. Soc. 104, p. 5849 (1982)), are known and will be apparent to those of ordinary skill in the art.

Compounds of formulae II, III and IV, and functionality-protected derivatives thereof, are useful as intermediates for the preparation of compounds of formula I. In those cases where L represents D', compounds of formula III may be converted to compounds of formula I by reaction with sulfonyl-activated species to form sulfonamides, sulfonyl ureas, thiocarbamates and the like. Methods for preparing such sulfonyl-activated species are well within the ordinary skill of the art. Typically, sulfonyl halides are used to obtain sulfonamides. Many sulfonyl halides


- 46 -

are commercially available; others may be easily obtained using conventional synthetic techniques (Gilbert, E.E. "Recent Developments in Preparative Sulfonation and Sulfation" Synthesis 1969: 3 (1969) and references cited therein; Hoffman, R.V. "M-Trifluoromethylbenzenesulfonyl Chloride" Org. Synth. Coll. Vol. VII, John Wiley and Sons (1990); Hartman, G.D. et. al. "4-Substituted Thiophene-and Furan-2-sulfonamides as Topical Carbonic Anhydrase Inhibitors" J. Med. Chem., 35, p. 3822 (1992) and references cited therein. Sulfonyl ureas are usually obtained by the reaction of an amine with sulfonyl chloride or a suitable equivalent such as sulfonyl-bis-imidazole or sulfonyl-bis-N-methyl imidazole. Thiocarbamates are typically obtained by the reaction of an alcohol with sulfonyl chloride or a suitable equivalent such as sulfonyl-bis-imidazole or sulfonyl-bis-N-methyl imidazole.

In the case of compounds of formula III wherein L is hydrogen, conversion of the resultant primary amine to a secondary amine may be carried out by known techniques. Such techniques include reaction with an alkyl halide or alkyl sulfonate, or by reductive alkylation with an aldehyde or carboxylic acid or activated derivative thereof using, for instance, catalytic hydrogenation or sodium cyanoborohydride (Borch et al., J. Am. Chem. Soc., 93, p. 2897 (1971)). Alternatively, the primary amine may be acylated followed by reduction with borane or another suitable reducing reagent, for example, as described by Cushman et al., J. Org. Chem., 56, p. 4161 (1991). This technique is especially useful in compounds of formula III where W represents a protecting group such as tert-butoxycarbonyl (Boc) or benzyloxycarbonyl (Cbz) and Q is H or where both W and Q are benzyl.



- 47 -




5

If variables W and Q of a particular compound of formula V represent removable protecting groups, removal of either or both groups followed by reaction of the resulting amine with an appropriate activated reagent will advantageously yield a different compound of formula V. For instance, reaction with an activated carboxylate, such as an acyl halide (e.g., acid fluorides, acid chlorides, and acid bromides), an activated ester such as 2- or 4-nitrophenyl esters, haloaryl esters (e.g., pentafluorophenyl or pentachlorophenyl) or 1-hydroxysuccinimide (HOSu) ester, a carbodiimide activated species, an anhydride, such as a symmetrical anhydride (e.g., isobutyl anhydride), or mixed carbonic-phosphoric or carbonic-phosphinic anhydrides, will yield the corresponding amide. Ureas may be obtained by reaction with isocyanates or amines in the presence of bis-activated carbonic acid derivatives such as phosgene or carbonyldiimidazole ("CDI"). Carbamates may be obtained by reaction with chlorocarbonates, with carbonates esterified with leaving groups such as 1-hydroxybenzotriazole ("HOBT"), HOSu, or 4-nitrophenol or with alcohols in the presence of bis-activated carbonic acid derivatives such as phosgene or its synthetic equivalents including diphosgene and triphosgene, or carbonyldiimidazole. An example of such a carbonate is N-succinimidyl-(3S)-tetrahydrofuran-3-yl carbonate. It will be readily recognized that in order to facilitate specific reactions, the protection of one or more potentially

- 48 -

reactive groups followed by subsequent removal of that group may be required. Such modification to the reaction schemes outlined above are within the ordinary skill of the art.

- 5           A particularly useful synthetic scheme for producing preferred sulfonamide intermediates of formula VIII is shown below wherein for compounds of formulas VI, VII and VIII, W and Q are as defined above for compounds of formula II, D' and E are as defined for compounds of formula I, and P' is H or amino protecting groups:
- 10



- 49 -

Compounds of formula VIII may be advantageously synthesized from readily available starting materials such as epoxide VI (see D.P. Getman, J. Med. Chem., 36, p. 288 (1993) and B.E. Evans et al., J. Org. Chem., 50, p. 4615 (1985)). Each step of the above synthetic scheme may be carried out as generally described above.

As can be appreciated by the skilled artisan, the above synthetic schemes are not intended to comprise a comprehensive list of all means by which the compounds described and claimed in this application may be synthesized. Further methods will be evident to those of ordinary skill in the art. Additionally, the various synthetic steps described above may be performed in an alternate sequence or order to give the desired compounds.

The compounds of this invention may be modified by appending appropriate functionalites to enhance selective biological properties. Such modifications are known in the art and include those which increase biological penetration into a given biological compartment (e.g., blood, lymphatic system, central nervous system), increase oral availability, increase solubility to allow administration by injection, alter metabolism and alter rate of excretion.

The novel compounds of the present invention are excellent ligands for aspartyl proteases, particularly HIV-1 and HIV-2 proteases. Accordingly, these compounds are capable of targeting and inhibiting late stage events in HIV replication, i.e., the processing of the viral polyproteins by HIV encoded proteases. Such compounds inhibit the proteolytic processing of viral polyprotein precursors by inhibiting aspartyl protease. Because aspartyl

- 50 -

protease is essential for the production of mature virions, inhibition of that processing effectively blocks the spread of virus by inhibiting the production of infectious virions, particularly from chronically infected cells. Compounds according to this invention advantageously inhibit the ability of the HIV-1 virus to infect immortalized human T cells over a period of days, as determined by an assay of extracellular p24 antigen -- a specific marker of viral replication. Other anti-viral assays have confirmed the potency of these compounds.

The compounds of this invention may be employed in a conventional manner for the treatment of viruses, such as HIV and HTLV, which depend on aspartyl proteases for obligatory events in their life cycle. Such methods of treatment, their dosage levels and requirements may be selected by those of ordinary skill in the art from available methods and techniques. For example, a compound of this invention may be combined with a pharmaceutically acceptable adjuvant for administration to a virally-infected patient in a pharmaceutically acceptable manner and in an amount effective to lessen the severity of the viral infection or to alleviate pathological effects associated with HIV infection or immunosuppression such as opportunistic infections or various cancers.

Alternatively, the compounds of this invention may be used in prophylactics and methods for protecting individuals against viral infection during a specific event, such as childbirth, or over an extended period of time. The compounds may be employed in such prophylactics either alone or together with other antiretroviral agents to enhance the efficacy of each agent. As such, the novel protease inhibitors of this

- 51 -

invention can be administered as agents for treating or preventing HIV infection in a mammal.

The compounds of formula I, especially those having a molecular weight of less than about 700 g/mole, may be readily absorbed into the bloodstream of mammals upon oral administration. Compounds of formula I having a molecular weight of less than about 600 g/mole and aqueous solubility of greater than or equal to 0.1 mg/mL are most likely to demonstrate high and consistent oral availability. This surprisingly impressive oral availability makes such compounds excellent agents for orally-administered treatment and prevention regimens against HIV infection.

In addition to being orally bioavailable, the compounds of this invention also have an impressively high therapeutic index (which measures toxicity versus anti-viral effect). Accordingly, the compounds of this invention are effective at lower dosage levels than many previously described conventional antiretroviral agents and avoid many of the severe toxic effects associated with those drugs. The potential of these compounds to be delivered at doses far exceeding their effective antiviral levels is advantageous in slowing or preventing the possibility of resistant variants developing.

The compounds of this invention may be administered to a healthy or HIV-infected patient either as a single agent or in combination with other anti-viral agents which interfere with the replication cycle of HIV. By administering the compounds of this invention with other anti-viral agents which target different events in the viral life cycle, the therapeutic effect of these compounds is potentiated. For instance, the co-administered anti-viral agent can be one which targets early events in the life cycle of

- 52 -

the virus, such as cell entry, reverse transcription and viral DNA integration into cellular DNA. Anti-HIV agents targeting such early life cycle events include, didanosine (ddI), dideoxycytidine (ddC), d4T, zidovudine (AZT), 3TC, 935U83, 1592U89, 524W91, polysulfated polysaccharides, sT4 (soluble CD4), ganciclovir, trisodium phosphonoformate, eflornithine, ribavirin, acyclovir, alpha interferon and trimetrexate. Additionally, non-nucleoside inhibitors of reverse transcriptase, such as TIBO, delavirdine (U90) or nevirapine, may be used to potentiate the effect of the compounds of this invention, as may viral uncoating inhibitors, inhibitors of trans-activating proteins such as tat or rev, or inhibitors of the viral integrase.

Combination therapies according to this invention exert an additive or synergistic effect in inhibiting HIV replication because each component agent of the combination acts on a different site of HIV replication. The use of such combination therapies may also advantageously reduce the dosage of a given conventional anti-retroviral agent which would be required for a desired therapeutic or prophylactic effect, as compared to when that agent is administered as a monotherapy. Such combinations may reduce or eliminate the side effects of conventional single anti-retroviral agent therapies, while not interfering with the anti-retroviral activity of those agents. These combinations reduce potential of resistance to single agent therapies, while minimizing any associated toxicity. These combinations may also increase the efficacy of the conventional agent without increasing the associated toxicity. In particular, we have discovered that in combination with other anti-HIV agents, the compounds of this invention act in an

- 53 -

additive or synergistical manner in preventing the replication of HIV in human T cells. Preferred combination therapies include the administration of a compound of this invention with AZT, ddI, ddC, d4T, 3TC, 935U83, 1592U89, 524W91 or a combination thereof.

Alternatively, the compounds of this invention may also be co-administered with other HIV protease inhibitors such as saquinavir (Ro 31-8959, Roche), MK 639 (Merck), ABT 538 (A-80538, Abbott), AG 1343 (Agouron), XM 412 (DuPont Merck), XM 450 (DuPont Merck), BMS 186318 (Bristol-Meyers Squibb) and CPG 53,437 (Ciba Geigy) or prodrugs of these or related compounds to increase the effect of therapy or prophylaxis against various viral mutants or members of HIV quasi species.

We prefer administering the compounds of this invention as single agents or in combination with retroviral reverse transcriptase inhibitors, such as nucleoside derivatives, or other HIV aspartyl protease inhibitors, including multiple combinations comprising from 3-5 agents. We believe that the co-administration of the compounds of this invention with retroviral reverse transcriptase inhibitors or HIV aspartyl protease inhibitors may exert a substantial additive or synergistic effect, thereby preventing, substantially reducing, or completely eliminating viral replication or infection or both, and symptoms associated therewith. Additionally, as the viruses are capable of developing resistance to certain aspartyl protease inhibitors quite rapidly, we believe that administration of a combination of agents may aid in slowing the development of resistant viruses relative to single agents alone.

The compounds of this invention can also be administered in combination with immunomodulators and

- 54 -

immunostimulators (e.g., bropirimine, anti-human alpha  
interferon antibody, IL-2, GM-CSF, interferon alpha,  
diethyldithiocarbamate, tumor necrosis factor,  
naltrexone, tuscarasol, and rEPO); and antibiotics  
5 (e.g., pentamidine isethionate) to prevent or combat  
infection and disease associated with HIV infections,  
such as AIDS, ARC and HIV-associated cancers.

When the compounds of this invention are  
administered in combination therapies with other  
10 agents, they may be administered sequentially or  
concurrently to the patient. Alternatively,  
pharmaceutical compositions according to this invention  
may comprise a combination of an aspartyl protease  
inhibitor of this invention and one or more therapeutic  
15 or prophylactic agents.

Although this invention focuses on the use of  
the compounds disclosed herein for preventing and  
treating HIV infection, the compounds of this invention  
can also be used as inhibitory agents for other viruses  
20 which depend on similar aspartyl proteases for  
obligatory events in their life cycle. These viruses  
include other AIDS-like diseases caused by  
retroviruses, such as simian immunodeficiency viruses,  
HTLV-I and HTLV-II. In addition, the compounds of this  
25 invention may also be used to inhibit other aspartyl  
proteases, and in particular, other human aspartyl  
proteases, including renin and aspartyl proteases that  
process endothelin precursors.

Pharmaceutical compositions of this invention  
30 comprise any of the compounds of the present invention,  
and pharmaceutically acceptable salts thereof, with any  
pharmaceutically acceptable carrier, adjuvant or  
vehicle. Pharmaceutically acceptable carriers,  
adjuvants and vehicles that may be used in the  
35 pharmaceutical compositions of this invention include,



- 55 -

but are not limited to, ion exchangers, alumina, aluminum stearate, lecithin, self-emulsifying drug delivery systems (SEDDS) such as  $\alpha$ -tocopherol polyethyleneglycol 1000 succinate, or other similar polymeric delivery matrices, serum proteins, such as human serum albumin, buffer substances such as phosphates, glycine, sorbic acid, potassium sorbate, partial glyceride mixtures of saturated vegetable fatty acids, water, salts or electrolytes, such as protamine sulfate, disodium hydrogen phosphate, potassium hydrogen phosphate, sodium chloride, zinc salts, colloidal silica, magnesium trisilicate, polyvinyl pyrrolidone, cellulose-based substances, polyethylene glycol, sodium carboxymethylcellulose, polyacrylates, waxes, polyethylene-polyoxypropylene-block polymers, polyethylene glycol and wool fat. Cyclodextrins such as  $\alpha$ -,  $\beta$ -, and  $\gamma$ -cyclodextrin, or chemically modified derivatives such as hydroxyalkylcyclodextrins, including 2- and 3-hydroxypropyl- $\beta$ -cyclodextrins, or other solublized derivatives may also be advantageously used to enhance delivery of compounds of formula I.

The pharmaceutical compositions of this invention may be administered orally, parenterally, by inhalation spray, topically, rectally, nasally, buccally, vaginally or via an implanted reservoir. We prefer oral administration or administration by injection. The pharmaceutical compositions of this invention may contain any conventional non-toxic pharmaceutically-acceptable carriers, adjuvants or vehicles. In some cases, the pH of the formulation may be adjusted with pharmaceutically acceptable acids, bases or buffers to enhance the stability of the formulated compound or its delivery form. The term parenteral as used herein includes subcutaneous, intracutaneous, intravenous, intramuscular, intra-

- 56 -

articular, intrasynovial, intrasternal, intrathecal, intralesional and intracranial injection or infusion techniques.

5 The pharmaceutical compositions may be in the form of a sterile injectable preparation, for example, as a sterile injectable aqueous or oleaginous suspension. This suspension may be formulated according to techniques known in the art using suitable dispersing or wetting agents (such as, for example, 10 Tween 80) and suspending agents. The sterile injectable preparation may also be a sterile injectable solution or suspension in a non-toxic parenterally-acceptable diluent or solvent, for example, as a solution in 1,3-butanediol. Among the acceptable 15 vehicles and solvents that may be employed are mannitol, water, Ringer's solution and isotonic sodium chloride solution. In addition, sterile, fixed oils are conventionally employed as a solvent or suspending medium. For this purpose, any bland fixed oil may be 20 employed including synthetic mono- or diglycerides. Fatty acids, such as oleic acid and its glyceride derivatives are useful in the preparation of injectables, as are natural pharmaceutically-acceptable oils, such as olive oil or castor oil, especially in 25 their polyoxyethylated versions. These oil solutions or suspensions may also contain a long-chain alcohol diluent or dispersant such as Ph. Hely or a similar alcohol.

30 The pharmaceutical compositions of this invention may be orally administered in any orally acceptable dosage form including, but not limited to, capsules, tablets, and aqueous suspensions and solutions. In the case of tablets for oral use, carriers which are commonly used include lactose and 35 corn starch. Lubricating agents, such as magnesium

- 57 -

stearate, are also typically added. For oral administration in a capsule form, useful diluents include lactose and dried corn starch. When aqueous suspensions are administered orally, the active ingredient is combined with emulsifying and suspending agents. If desired, certain sweetening and/or flavoring and/or coloring agents may be added.

The pharmaceutical compositions of this invention may also be administered in the form of suppositories for rectal administration. These compositions can be prepared by mixing a compound of this invention with a suitable non-irritating excipient which is solid at room temperature but liquid at the rectal temperature and therefore will melt in the rectum to release the active components. Such materials include, but are not limited to, cocoa butter, beeswax and polyethylene glycols.

Topical administration of the pharmaceutical compositions of this invention is especially useful when the desired treatment involves areas or organs readily accessible by topical application. For application topically to the skin, the pharmaceutical composition should be formulated with a suitable ointment containing the active components suspended or dissolved in a carrier. Carriers for topical administration of the compounds of this invention include, but are not limited to, mineral oil, liquid petroleum, white petroleum, propylene glycol, polyoxyethylene polyoxypropylene compound, emulsifying wax and water. Alternatively, the pharmaceutical composition can be formulated with a suitable lotion or cream containing the active compound suspended or dissolved in a carrier. Suitable carriers include, but are not limited to, mineral oil, sorbitan monostearate, polysorbate 60, cetyl esters wax, cetearyl alcohol,

- 58 -

2-octyldodecanol, benzyl alcohol and water. The pharmaceutical compositions of this invention may also be topically applied to the lower intestinal tract by rectal suppository formulation or in a suitable enema formulation. Topically-transdermal patches are also included in this invention.

The pharmaceutical compositions of this invention may be administered by nasal aerosol or inhalation. Such compositions are prepared according to techniques well-known in the art of pharmaceutical formulation and may be prepared as solutions in saline, employing benzyl alcohol or other suitable preservatives, absorption promoters to enhance bioavailability, fluorocarbons, and/or other solubilizing or dispersing agents known in the art.

Dosage levels of between about 0.01 and about 100 mg/kg body weight per day, preferably between about 0.5 and about 75 mg/kg body weight per day of the active ingredient compound are useful in the prevention and treatment of viral infection, including HIV infection. Typically, the pharmaceutical compositions of this invention will be administered from about 1 to about 5 times per day or alternatively, as a continuous infusion. Such administration can be used as a chronic or acute therapy. The amount of active ingredient that may be combined with the carrier materials to produce a single dosage form will vary depending upon the host treated and the particular mode of administration. A typical preparation will contain from about 5% to about 95% active compound (w/w). Preferably, such preparations contain from about 20% to about 80% active compound.

Upon improvement of a patient's condition, a maintenance dose of a compound, composition or combination of this invention may be administered, if necessary. Subsequently, the dosage or frequency of

- 59 -

administration, or both, may be reduced, as a function of the symptoms, to a level at which the improved condition is retained when the symptoms have been alleviated to the desired level, treatment should  
5 cease. Patients may, however, require intermittent treatment on a long-term basis upon any recurrence of disease symptoms.

As the skilled artisan will appreciate, lower or higher doses than those recited above may be  
10 required. Specific dosage and treatment regimens for any particular patient will depend upon a variety of factors, including the activity of the specific compound employed, the age, body weight, general health status, sex, diet, time of administration, rate of  
15 excretion, drug combination, the severity and course of the infection, the patient's disposition to the infection and the judgment of the treating physician.

The compounds of this invention are also useful as commercial reagents which effectively bind to  
20 aspartyl proteases, particularly HIV aspartyl protease. As commercial reagents, the compounds of this invention, and their derivatives, may be used to block proteolysis of a target peptide or may be derivatized to bind to a stable resin as a tethered substrate for  
25 affinity chromatography applications. For example, a compound of formula I may be tethered to an affinity column to purify recombinantly produced HIV protease. Derivatization of the compounds of this invention to  
30 produce affinity chromatography resins and the methods used to purify proteases using such resins are well known and within the skill of the art. These and other uses which characterize commercial aspartyl protease inhibitors will be evident to those of ordinary skill in the art. (See: Rittenhouse, J. et al. Biochem.

- 60 -

Biophys. Res. Commun. 171, p. 60 (1990) and Heimbach, J.C. et al. Ibid 164, p. 955 (1989)).

In order that this invention be more fully understood, the following examples are set forth.

5 These examples are for the purpose of illustration only and are not to be construed as limiting the scope of the invention in any way.

#### General Materials and Methods

10 All temperatures are recorded in degrees Celsius. Thin layer chromatography (TLC) was carried out using 0.25 mm thick E. Merck silica gel 60 F<sub>254</sub> plates and elution with the indicated solvent system. Detection of the compounds was carried out by treating the plate with an appropriate visualizing agent, such as 10% solution of phosphomolybdic acid in ethanol or a 0.1% solution of ninhydrin in ethanol, followed by heating, and/or by exposure to UV light or iodine vapors when appropriate. Thick layer silica gel chromatography was also carried out using E. Merck 60 F<sub>254</sub> plates ("prep plates") of 0.5, 1.0, or 2.0 mm thickness. Following development of the plate, the band of silica containing the desired compound was isolated and eluted with an appropriate solvent. Analytical HPLC was carried out using a Water's Delta 25 Pak, 5  $\mu$ M silica, C18 reversed-phase column, 3.9 mm ID x 15 cm L with a flow rate of 1.5 mL/min using the following table:

	Mobile phase:	A = 0.1% CF <sub>3</sub> CO <sub>2</sub> H in H <sub>2</sub> O
		B = 0.1% CF <sub>3</sub> CO <sub>2</sub> H in CH <sub>3</sub> CN
30	Gradient:	T = 0 min., A (95%), B (5%)
		T = 20 min., A (0%), B (100%)
		T = 22.5 min., A (0%), B (100%)

- 61 -

Preparative HPLC was also carried out using C<sub>18</sub> reversed-phase media. HPLC retention times were recorded in minutes. NMR spectral data was recorded using a Bruker AMX500, equipped with either a reverse or QNP probe, at 500 MHz, and was taken in the indicated solvent.

We have measured the inhibition constants of each compound against HIV-1 protease using the method described essentially by M.W. Pennington et al., Peptides 1990, Gimet, E. and D. Andrew, Eds., Escom; Leiden, Netherlands (1990).

Compounds of formula I were tested for their antiviral potency in several virological assays. In the first assay, the compounds were added as a solution in dimethylsulfoxide (DMSO) to a test cell culture of CCRM-CEM cells, a strain of CD4<sup>+</sup> human T-cell lymphoma cells, previously acutely infected with HIV<sub>1</sub> using standard protocols (see Meek, T. D. et al., "Inhibition of HIV-1 protease in infected T-lymphocytes by synthetic peptide analogues", Nature, 343, p. 90 (1990)). Preferred compounds are those which are able to inhibit 90% of viral infectivity at a concentration of 1  $\mu$ M or less. More preferred compounds are those which are able to inhibit 90% of viral infectivity at a concentration of 100 nM or less.

The effect of the compounds on inhibiting the replication of the virus was measured by determining the HIV extracellular p24 antigen concentration using a commercial enzyme immunoassay (obtained from Coulter Corporation, Hialeah, FL).

Depending on the cell type and the desired readout, syncytia formation, reverse-transcriptase (RT) activity, or cytopathic effect as assayed by a dye uptake method may also be used as readouts of antiviral activity. See H. Mitsuya and S. Broder, "Inhibition of

- 62 -

the in vitro infectivity and cytopathic effect of human T-lymphotropic virus type III/lymphadenopathy-associated virus (HTLV-III/LAV) by 2',3'-dideoxynucleosides", Proc. Natl. Acad. Sci. USA, vol. 83, pp. 1911-1915 (1986). The effect of compounds of formula I on clinical isolates of other HIV-1 strains was determined by obtaining low-passaged virus from HIV-infected patients and assaying the effect of the inhibitors in preventing infection of the HIV virus in freshly prepared human peripheral blood mononuclear cells (PBMCs).

Insofar as compounds of formula I are able to inhibit the replication of the HIV virus in human T-cells and furthermore, may be delivered orally to mammals, they are of evident clinical utility for the treatment of HIV infection. These tests are predictive of the compounds ability to inhibit HIV protease in vivo.

#### Example 1

##### Synthesis of Compound 35

A. Compound VII (D' = isobutyl, W = tert-butoxy carbonyl, Q = H, P' = H). A solution of 4.1 g of epoxide VI (W=Boc, Q=H) in 30 mL of ethanol was treated with 22.4 mL of isobutylamine and heated under reflux for 1 h. The mixture was concentrated to yield the title compound as a white solid which was used without subsequent purification. NMR (CDCl<sub>3</sub>):  $\delta$  0.91 (d, 3H); 0.93 (d, 3H); 1.37 (s, 9H); 1.68 (br s, 2H); 2.40 (d, 2H); 2.68 (d, 2H); 2.87 (dd, 1H); 2.99 (dd, 1H); 3.46 (dd, 1H); 3.75 (br s, 1H); 3.80 (br s, 1H); 4.69 (d, 1H); 7.19-7.32 (m, 4H).

B. Compound 32. A solution of 391 mg of the resultant compound of Example 1A in 4:1 CH<sub>2</sub>Cl<sub>2</sub>/saturated aqueous



- 63 -

NaHCO<sub>3</sub> was treated sequentially, at ambient temperature under an atmosphere of nitrogen, with 271 mg of 4-fluorobenzenesulfonyl chloride and 117 mg of sodium bicarbonate. The mixture was stirred for 14 h, diluted with CH<sub>2</sub>Cl<sub>2</sub>, washed with saturated NaCl then dried over MgSO<sub>4</sub>, filtered, and concentrated in vacuo. The residue was purified by low pressure silica gel chromatography using 5% diethyl ether in CH<sub>2</sub>Cl<sub>2</sub> as eluent to yield 420 mg of the title compound as a white solid. TLC: R<sub>f</sub> = 0.20, 5% diethyl ether in CH<sub>2</sub>Cl<sub>2</sub>. HPLC: R<sub>t</sub> = 17.41 min; (<sup>1</sup>H)-NMR (CDCl<sub>3</sub>) consistent with structure.

C. Compound VIII (W = H, Q = H, D' = isobutyl, E = 4-fluorophenyl, hydrochloride salt). A solution of 398 mg of the resultant compound of Example 1B in ethyl acetate was treated at -20°C with HCl gas. The HCl was bubbled through the mixture for 20 min over which time the temperature was allowed to warm to 20°C. Nitrogen was then bubbled through the mixture for 15 min and solvent removed in vacuo to yield 347 mg of the title compound as a white solid. TLC: R<sub>f</sub> = 0.82, 5:10:85 NH<sub>4</sub>OH/CH<sub>3</sub>OH/CH<sub>2</sub>Cl<sub>2</sub>; (<sup>1</sup>H)-NMR (CDCl<sub>3</sub>) consistent with structure.

D. Compound 35. A solution of 111 mg of the resultant compound of Example 1C in CH<sub>2</sub>Cl<sub>2</sub> was added, at ambient temperature under an atmosphere of nitrogen, to a solution of 118 mg of N-succinimidyl-(S)-3-tetrahydrofuranyl carbonate (hereinafter "THF-OSu") and 133 mg N,N-diisopropylethylamine in CH<sub>2</sub>Cl<sub>2</sub>. The mixture was stirred for 14 h, diluted with CH<sub>2</sub>Cl<sub>2</sub>, washed with saturated NaHCO<sub>3</sub> and saturated NaCl, then dried over MgSO<sub>4</sub>, filtered, and concentrated in vacuo. The

- 64 -

residue was subjected to preparative thin layer silica gel chromatography using 5% CH<sub>3</sub>OH in CH<sub>2</sub>Cl<sub>2</sub> to yield 98.8 mg of the title compound as a white solid. TLC: R<sub>f</sub> = 0.48, 5% CH<sub>3</sub>OH in CH<sub>2</sub>Cl<sub>2</sub>. HPLC: R<sub>t</sub> = 15.18 min; (1H)-NMR (CDCl<sub>3</sub>) consistent with structure.

### Example 2

#### Synthesis of Compound 101

A. Compound VII (W = tert-butoxycarbonyl, Q = H, D' = methyl, P' = H). To a solution of compound VI (W=Boc, Q=H) (1.7 mmol) in ethanol (20 mL) was added methylamine gas, at ambient temperature, for 30 min. The solution was stirred overnight, then concentrated under reduced pressure to give 0.47 g of the title compound which was used without subsequent purification. TLC: R<sub>f</sub> = 0.19, 1:10:90 NH<sub>4</sub>OH/methanol/CH<sub>2</sub>Cl<sub>2</sub>, (1H)-NMR (CDCl<sub>3</sub>) consistent with structure.

B. Compound 128. To a solution of the product of Example 2A (0.15 g, 0.51 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (6 mL) was added a saturated solution of sodium bicarbonate (3 mL), followed by addition of solid sodium bicarbonate (90 mg, 1.0 mmol), followed by addition of acetamidobenzenesulfonyl chloride (0.24 g, 1.02 mmol). The mixture was stirred at ambient temperature overnight. The organics were extracted into 100 mL CH<sub>2</sub>Cl<sub>2</sub>, dried over anhydrous, MgSO<sub>4</sub>, concentrated under reduced pressure then purified via medium pressure silica gel chromatography using a gradient system of CH<sub>2</sub>Cl<sub>2</sub> followed by 5:95 EtOAc/CH<sub>2</sub>Cl<sub>2</sub>, followed by 10:90 EtOAc/CH<sub>2</sub>Cl<sub>2</sub>. The title compound was obtained as 244 mg of white solid. TLC: R<sub>f</sub> = 0.13, 3:97 methanol/CH<sub>2</sub>Cl<sub>2</sub>,

- 65 -

HPLC: Rt = 13.47 min; (<sup>1</sup>H)-NMR (CDCl<sub>3</sub>) consistent with structure.

5 C. Compound 101. This compound was prepared from the resultant compound of Example 2B by treatment with hydrogen chloride gas as described in Example 1C and subsequent reaction of this material with THF-OSu in the manner described in Example 1D. After workup and purification by preparative reversed-phase C<sub>18</sub> HPLC using a linear gradient of 35% to 100% CH<sub>3</sub>CN/H<sub>2</sub>O with 0.1% TFA as eluant on a portion of the crude mixture, 10 4.2 mg of the title compound was obtained as a white solid. TLC: Rf = 0.2, 4% MeOH/CH<sub>2</sub>Cl<sub>2</sub>. HPLC: Rt = 11.53 min; (<sup>1</sup>H)-NMR (CDCl<sub>3</sub>) consistent with structure.

### Example 3

#### 15 Synthesis of Compound 116

A. Aminomethylcyclopentane. To a solution of LiAlH<sub>4</sub> (38 g, 1.0 mole) in diethyl ether (2 L) was added cyclopentanecarbonitrile (73.2 g, 0.77 mol) as a solution in 250 mL ether. The solution was stirred overnight at ambient temperature and then quenched by addition of the organics to 3 L of a saturated potassium, sodium tartrate solution. The amine was extracted into 3 L of ether, dried over anhydrous K<sub>2</sub>CO<sub>3</sub>, then concentrated by distillation to approximately 400 20 mL total volume. The crude product was purified via distillation to give 58.2 g of the title compound as a colorless oil. (<sup>1</sup>H)-NMR (CDCl<sub>3</sub>) consistent with structure. 25

30 B. Compound VII (W = tert-butoxycarbonyl, Q = H, D' = cyclopentylmethyl, P' = H). To the resultant compound of Example 3A (20 g, 0.2 mol) was added compound VI (W

- 66 -

= Boc, Q = H) (5.84 g) and the mixture was stirred for 24 h at ambient temperature. The solution was concentrated by distillation under reduced pressure. The residue was triturated with hexane and the solid collected by suction filtration and washed with hexane to give 7.08 g of a white solid which was used without further purification. TLC: Rf = 0.59 (1:10:90 concentrated NH<sub>4</sub>OH/methanol/CH<sub>2</sub>Cl<sub>2</sub>), (<sup>1</sup>H)-NMR (CDCl<sub>3</sub>) consistent with structure.

10 C. Compound VIII (W = tert-butoxycarbonyl, Q = H, D' = cyclopentylmethyl, E = 4-chlorophenyl). The resultant compound of Example 3B (252 mg) was reacted with 4-chlorobenzenesulfonyl chloride (175 mg) in the manner described in Example 8H. Workup and  
15 purification by silica gel chromatography using EtOAc/CH<sub>2</sub>Cl<sub>2</sub> as eluant yielded the product as a white solid; (<sup>1</sup>H) NMR (CDCl<sub>3</sub>) consistent with structure.

D. Compound VIII (W = H, Q = H, D' = cyclopentylmethyl, E = 4-chlorophenyl, hydrochloride salt). A solution of 320 mg of the resultant compound of Example 3C in 20 mL of EtOAc was treated with  
20 anhydrous HCl gas for 5 min. The reaction mixture was purged with nitrogen then concentrated in vacuo to yield a white solid which was used directly for subsequent  
25 reaction.

E. Compound 116. To a solution of 63.4 mg of the resultant compound of Example 3D in 1 mL of THF was added sequentially 54 μL of diisopropylethylamine and a solution of 39.9 mg of THF-OSu in 1 mL THF. The  
30 mixture was stirred for 24 hours and then concentrated in vacuo. The residue was purified by low pressure silica gel column chromatography using 20% EtOAc in

- 67 -

CH<sub>2</sub>Cl<sub>2</sub> eluent to yield 0.62 g of the title compound.  
TLC: R<sub>f</sub> = 0.71, 40% EtOAc/CH<sub>2</sub>Cl<sub>2</sub>. HPLC: R<sub>t</sub> = 16.88 min.  
(<sup>1</sup>H) NMR (CDCl<sub>3</sub>) consistent with structure.

#### Example 4

5

#### Synthesis of Compound 132

A. Compound VII (W = tert-butoxycarbonyl, Q = H, D' = (2-tetrahydrofuryl)-methyl, P' = H). To a solution of compound VI (W=Boc, Q=H) (3.3 mmol) in ethanol (30 mL) was added tetrahydrofurfurylamine (1.03 mL, 10 mmol).  
10 The mixture was warmed to 85°C and stirred overnight. The solution was filtered and the solution concentrated under reduced pressure to give 1.29 g of the title compound which was used without subsequent purification. TLC: R<sub>f</sub> = 0.52, 1:10:90  
15 NH<sub>4</sub>OH/methanol/CH<sub>2</sub>Cl<sub>2</sub>,

B. Compound 129. To a solution of the resultant compound of Example 4A (200 mg, 0.55 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (6 mL) was added 4-fluorobenzenesulfonyl chloride (320 mg, 1.6 mmol) followed a saturated solution of sodium  
20 bicarbonate (3 mL) and solid sodium bicarbonate (0.1 g, 1.2 mmol). The mixture was stirred at ambient temperature overnight. The solution was diluted with 100 mL CH<sub>2</sub>Cl<sub>2</sub>, the organics separated, dried over anhydrous MgSO<sub>4</sub>, and the organics concentrated under  
25 reduced pressure. The crude product was purified via medium pressure liquid chromatography using a gradient solvent system of CH<sub>2</sub>Cl<sub>2</sub> followed by 5:95 ether/CH<sub>2</sub>Cl<sub>2</sub> followed by a 10:90 ether/CH<sub>2</sub>Cl<sub>2</sub> solution to give 130 mg of the title compound as a white solid. TLC: R<sub>f</sub> =  
30 0.35, 3:97 methanol/CH<sub>2</sub>Cl<sub>2</sub>, HPLC: R<sub>t</sub> = 16.37 min, (<sup>1</sup>H)-NMR (CDCl<sub>3</sub>) consistent with structure.

- 68 -

C. Compound VIII (W = H, Q = H, D' = (2-tetrahydrofuryl)-methyl, E = 4-fluorophenyl, hydrochloride salt). To a solution of the resultant compound of Example 4B (30 mg, 0.057 mmol) in EtOAc (3 mL) was added 30% w/w HCl in EtOAc (1 mL). The mixture was stirred overnight at ambient temperature. The solution was concentrated under reduced pressure to give 16 mg of the title compound as a white solid which was used without subsequent purification. TLC: Rf = 0.60 (1:10:90 NH<sub>4</sub>OH/methanol/CH<sub>2</sub>Cl<sub>2</sub>).

D. Compound 132. To a solution of the resultant compound of Example 4C (16 mg) in CH<sub>2</sub>Cl<sub>2</sub> (5 mL) was added triethylamine (0.1 mL, 0.72 mmol) followed by THF-OSu (20 mg, 0.09 mmol). The mixture was stirred at ambient temperature for 24 hours. The solution was concentrated under reduced pressure and the crude product purified via medium pressure column chromatography using 20:80 EtOAc/CH<sub>2</sub>Cl<sub>2</sub> as the solvent system to give 7.4 mg. Rf = 0.37 (3:97 methanol/CH<sub>2</sub>Cl<sub>2</sub>), HPLC: Rt = 14.19 min, (<sup>1</sup>H)-NMR (CDCl<sub>3</sub>) consistent with structure.

#### Example 5

##### Synthesis of Compound 134

A. Compound VII (W = tert-butoxycarbonyl, Q = H, D' = (isobutenyl, P' = H)). To a solution compound VI (W = tert-butoxycarbonyl, Q = H) (2.5 mmol) in ethanol (30 mL) was added a solution 2-methallylamine hydrochloride (1.34 g, 12.5 mmol) and KOH (0.70 g, 12.5 mmol) in ethanol (20 mL). The mixture stirred 30 min at ambient temperature. The solutions were combined and heated to 85°C for 24 h. The solution was filtered and concentrated under reduced pressure to give 0.82 g of the title compound which was used without subsequent

- 69 -

purification. TLC:  $R_f$  = 0.45, 1:10:90 concentrated  $\text{NH}_4\text{OH}$ /methanol/ $\text{CH}_2\text{Cl}_2$ .

5 B. Compound 131. To a solution of the resultant compound of Example 5A (200 mg, 0.60 mmol) in  $\text{CH}_2\text{Cl}_2$  (6 mL) was added 4-acetamidobenzenesulfonyl chloride (410 mg, 1.76 mmol), followed by a saturated solution of sodium bicarbonate (3 mL) and solid sodium bicarbonate (0.1 g, 1.2 mmol). The mixture was stirred at ambient temperature overnight. The solution was  
10 diluted with 100 mL  $\text{CH}_2\text{Cl}_2$ , the organics separated, dried over anhydrous  $\text{MgSO}_4$ , and the organics concentrated under reduced pressure. The crude product was purified via medium pressure liquid chromatography using a gradient solvent system of  $\text{CH}_2\text{Cl}_2$ , followed by  
15 30:70 EtOAc/ $\text{CH}_2\text{Cl}_2$  solution to give 140 mg of the title compound as a white solid. TLC:  $R_f$  = 0.19, 3:97 methanol/ $\text{CH}_2\text{Cl}_2$ , HPLC:  $R_t$  = 15.06 min, ( $^1\text{H}$ )-NMR ( $\text{CDCl}_3$ ) consistent with structure.

20 C. Compound VIII (W = H, Q = H, D' = isobutenyl, E = 4-acetamidophenyl, hydrochloride salt). To a solution of the resultant compound of Example 5B (40 mg 0.075 mmol) in EtOAc (5 mL) was added 30% w/w HCl in EtOAc (2 mL). The mixture was stirred overnight at ambient temperature. The solution was concentrated under  
25 reduced pressure to give the title compound, which was used without subsequent purification. TLC:  $R_f$  = 0.38, 1:10:90  $\text{NH}_4\text{OH}$ /methanol/ $\text{CH}_2\text{Cl}_2$ .

30 D. Compound 134. To a solution of the resultant compound of Example 5C in  $\text{CH}_2\text{Cl}_2$  (5 mL) was added triethylamine (0.1 mL, 0.72 mmol), followed by THF-OSu (26 mg, 0.11 mmol). The mixture was stirred at ambient temperature for 24 hours. The solution was

- 70 -

concentrated under reduced pressure and the crude product purified via medium pressure column chromatography using a gradient solvent system of  $\text{CH}_2\text{Cl}_2$ , followed by 1:99 methanol/ $\text{CH}_2\text{Cl}_2$ , followed by 3:97 methanol/ $\text{CH}_2\text{Cl}_2$ , as the solvent system to give 10.1 mg of the title compound.  $R_f = 0.11$  (3:97 methanol/ $\text{CH}_2\text{Cl}_2$ ), HPLC:  $R_t = 12.86$  min,  $(^1\text{H})$ -NMR ( $\text{CDCl}_3$ ) consistent with structure.

#### Example 6

10

#### Synthesis of Compound 136

A. Compound VII ( $W = \text{tert-butoxycarbonyl}$ ,  $Q = \text{H}$ ,  $D' = 2\text{-furfuryl}$ ,  $P' = \text{H}$ ). To a solution compound VI ( $W = \text{Boc}$ ,  $Q = \text{H}$ ) (2.5 mmol) in ethanol (30 mL) was added furfurylamine (0.67 mL, 7.5 mmol) and the mixture was heated to  $85^\circ\text{C}$  for 24 h. The solution was filtered and concentrated under reduced pressure to give 0.80 g of the title compound which was used without subsequent purification. TLC:  $R_f = 0.38$ , 1:10:90 concentrated  $\text{NH}_4\text{OH}/\text{methanol}/\text{CH}_2\text{Cl}_2$ .

20 B. Compound VIII ( $W = \text{tert-butoxycarbonyl}$ ,  $Q = \text{H}$ ,  $D' = 2\text{-furyl}$ ,  $E = 4\text{-fluorophenyl}$ ). To a solution of the product of Example 6A (0.20 g, 0.60 mmol) in  $\text{CH}_2\text{Cl}_2$  (6 mL) was added a saturated solution of sodium bicarbonate (3 mL), followed by addition of solid sodium bicarbonate (0.1 g, 1.2 mmol), then p-fluorobenzenesulfonyl chloride (0.32 g, 1.6 mmol). The mixture was stirred at ambient temperature for 24 h. The organics were extracted into 100 mL  $\text{CH}_2\text{Cl}_2$ , dried over anhydrous  $\text{MgSO}_4$ , concentrated under reduced pressure, then purified via medium pressure silica gel chromatography using a gradient system of  $\text{CH}_2\text{Cl}_2$ , followed by 1:99 methanol/ $\text{CH}_2\text{Cl}_2$ . The title compound was obtained as a white solid (86.1 mg). TLC:  $R_f =$



- 71 -

0.17, 3:97 methanol/ $\text{CH}_2\text{Cl}_2$ , HPLC:  $R_t$  = 16.5 min; ( $^1\text{H}$ )-NMR ( $\text{CDCl}_3$ ) consistent with structure.

5 C. Compound VIII ( $W = \text{H}$ ,  $Q = \text{H}$ ,  $D' = 2\text{-furyl}$ ,  $E = 4\text{-fluorophenyl}$ , hydrochloride salt). To a solution of the resultant compound of Example 6B (16 mg, 0.031 mmol) in EtOAc (3 mL) was added 30% w/w HCl in EtOAc (1 mL). The mixture was stirred overnight at ambient temperature. The solution was concentrated under reduced pressure to give the title compound, which was used without subsequent purification. TLC:  $R_f$  = 0.48, 1:10:90  $\text{NH}_4\text{OH}$ /methanol/ $\text{CH}_2\text{Cl}_2$ .

10 D. Compound 136. To a solution of the resultant compound of Example 6C in  $\text{CH}_2\text{Cl}_2$  (5 mL) was added triethylamine (0.1 mL, 0.72 mmol), followed by THF-OSu (11 mg, 0.05 mmol). The mixture was stirred at ambient temperature for 24 hours. The solution was concentrated under reduced pressure and the crude product purified via medium pressure column chromatography using a gradient solvent system of  $\text{CH}_2\text{Cl}_2$ , followed by 20:80 EtOAc/ $\text{CH}_2\text{Cl}_2$ , as the solvent system to give 4.9 mg. TLC:  $R_f$  = 0.28, (3:97 methanol/ $\text{CH}_2\text{Cl}_2$ , HPLC:  $R_t$  = 14.57 min, ( $^1\text{H}$ )-NMR ( $\text{CDCl}_3$ ) consistent with structure.

#### Example 7

##### 25 Synthesis of Compound 158

A. Compound VII ( $W = \text{tert-butoxycarbonyl}$ ,  $Q = \text{H}$ ,  $D' = \text{cyclohexylmethyl}$ ,  $P' = \text{H}$ ). To a solution of compound VI ( $W = \text{Boc}$ ,  $Q = \text{H}$ ) (5.0 mmol) in ethanol (20 mL) was added cyclohexylmethanamine (3.25 mL, 2.83 mmol) and the mixture was stirred for 3 hours at ambient temperature. The solution was filtered and

30

- 72 -

concentrated under reduced pressure to give 1.49 g of a white solid which was used directly for subsequent reaction. TLC:  $R_f = 0.14$ , 3:97 methanol/ $\text{CH}_2\text{Cl}_2$ . ( $^1\text{H}$ )-NMR ( $\text{CDCl}_3$ ) consistent with structure.

- 5 B. Compound VIII ( $W = \text{tert-butoxycarbonyl}$ ,  $Q = \text{H}$ ,  $D' = \text{cyclohexylmethyl}$ ,  $E = 4\text{-methoxyphenyl}$ ). To a solution of the resultant compound of Example 7A (400 mg, 1.06 mmol) in  $\text{CH}_2\text{Cl}_2$  (10 mL) was added 4-methoxybenzenesulfonyl chloride (0.66 g, 3.1 mmol)
- 10 followed by addition of a saturated solution of sodium bicarbonate (3 mL) and 0.18 g of solid sodium bicarbonate. The mixture was stirred at ambient temperature overnight. The solution was diluted with 200 mL  $\text{CH}_2\text{Cl}_2$ , the organics separated, dried over
- 15 anhydrous  $\text{MgSO}_4$ , and the organics concentrated under reduced pressure. The crude product was purified via medium pressure liquid chromatography using  $\text{CH}_2\text{Cl}_2$ , followed by 1:99 methanol/ $\text{CH}_2\text{Cl}_2$  as the solvent system to give 340 mg of the title compound as a white solid.
- 20 TLC:  $R_f = 0.39$ , 3:97 methanol/ $\text{CH}_2\text{Cl}_2$ , ( $^1\text{H}$ )-NMR ( $\text{CDCl}_3$ ) consistent with structure.

- C. Compound VIII ( $W = \text{H}$ ,  $Q = \text{H}$ ,  $D' = \text{cyclohexylmethyl}$ ,  $E = 4\text{-methoxyphenyl}$ , hydrochloride salt). To a solution of the resultant compound of
- 25 Example 7B (0.34 g, 0.62 mmol) in EtOAc (10 mL) was added 30% w/w HCl in EtOAc (5 mL). The mixture was stirred for 3 hours at ambient temperature. The solution was concentrated under reduced pressure to give 0.3 g of a white solid which was used directly for
- 30 subsequent reaction. TLC:  $R_f = 0.12$ , 3:97 methanol/ $\text{CH}_2\text{Cl}_2$ .

- 73 -

D. Compound 158. To a solution of the resultant compound of Example 7C (100 mg, 0.21 mmol) in  $\text{CH}_2\text{Cl}_2$  (8 mL) was added triethylamine (0.2 mL, 1.44 mmol) followed by THF-OSu (71 mg, 0.31 mmol). The mixture was stirred at ambient temperature for 6 hours. The solution was diluted with  $\text{CH}_2\text{Cl}_2$ , (200 mL) washed with a saturated solution of sodium bicarbonate (30 mL), the organics separated, dried over anhydrous  $\text{MgSO}_4$  and concentrated under reduced pressure and the crude product purified via medium pressure column chromatography using a gradient solvent system of  $\text{CH}_2\text{Cl}_2$  followed by 10:90 EtOAc/ $\text{CH}_2\text{Cl}_2$  as the solvent system to give 84.9 mg of the title compound. TLC:  $R_f$  = 0.48, 3:97 methanol/ $\text{CH}_2\text{Cl}_2$ , HPLC:  $R_t$  = 16.35 min; ( $^1\text{H}$ )-NMR ( $\text{CDCl}_3$ ) consistent with structure.

#### Example 8

##### Synthesis of Compound 195

A. 3(S)-amino-2(syn)-hydroxy-4-phenyl-1-chlorobutane formate salt. To a slurry of 16.33 g of 10% palladium on carbon (25% by weight) in methanol and tetrahydrofuran (400 mL, 1:1) was added, under  $\text{N}_2$ , 65.35 g of 3(S)-N-(-benzyloxycarbonyl)-amino-1-chloro-2(syn)-hydroxy-4-phenylbutane (195.77 mmol) as a solution in methanol and tetrahydrofuran (1.2 L). To this slurry was added 540 mL of formic acid. After 15 h, the reaction mixture was filtered through diatomaceous earth and concentrated to dryness. The resultant oil was slurried in toluene and evaporated, then triturated sequentially with diethyl ether and  $\text{CH}_2\text{Cl}_2$  to provide 47.64 g of product as a granular tan solid. TLC:  $R_f$  = 0.17, 5% acetic acid/ethyl acetate.

- 74 -

B. 3(S)-N-(3(S)-tetrahydrofuryloxycarbonyl)-amino-1-chloro-2(syn)-hydroxy-4-phenylbutane. To a solution of the resultant compound of Example 8A (1.97 g, 7.95 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (20 mL) was added a saturated solution of sodium bicarbonate (5 mL), followed by solid sodium bicarbonate (1.33 g, 17.9 mmol), and THF-OSu (2.0 g, 8.7 mmol). The mixture was stirred at ambient temperature overnight. The solution was diluted with 200 mL CH<sub>2</sub>Cl<sub>2</sub>, the organics separated, dried over anhydrous MgSO<sub>4</sub>, and concentrated under reduced pressure. The residue was recrystallized from ethyl acetate/hexane to give 1.01 g of the title compound as a white solid. TLC: R<sub>f</sub> = 0.35, 3:97 methanol/CH<sub>2</sub>Cl<sub>2</sub>. (<sup>1</sup>H)-NMR (CDCl<sub>3</sub>) consistent with structure.

C. Compound VI (W = H, Q = 3(s)-tetrahydrofuryloxycarbonyl). To a solution of the resultant compound of Example 8B (1.0 g, 3.2 mmol) in absolute ethanol (15 mL) was added solid KOH (0.21 g, 3.8 mmol). The mixture was stirred at ambient temperature for 1.0 h. The solution was filtered through a pad of Celite then concentrated under reduced pressure. The residue was taken up in ether (100 mL), washed with brine, dried over MgSO<sub>4</sub>, the concentrated under reduced pressure to give 0.88 g of the title compound as a white solid. TLC: R<sub>f</sub> = 0.49 (3:97 methanol/CH<sub>2</sub>Cl<sub>2</sub>), (<sup>1</sup>H)-NMR (CDCl<sub>3</sub>) consistent with structure.

D. Compound III (W = H, Q = (S)-3-tetrahydrofuryloxycarbonyl, D = benzyl, D' = cyclopentylmethyl, R, = H, L = H). The resultant compound of Example 8C (0.88 g, 3.2 mol) was added to the resultant compound of Example 3A (5.0 g, 50.4 mmol) and stirred for 24 h at ambient temperature. The

- 75 -

solution was concentrated by distillation under reduced pressure. The residue was triturated with hexane and the solid collected by suction filtration and washed with hexane to give 0.93 g of the title compound. TLC: R<sub>f</sub> = 0.44, 1:10:90 concentrated NH<sub>4</sub>OH/methanol/CH<sub>2</sub>Cl<sub>2</sub>; (H)-NMR (CDCl<sub>3</sub>) consistent with structure.

E. Compound VII (W = H, Q = (S)-3-tetrahydrofuryl, D' = cyclopentylmethyl, P' = tert-butoxycarbonyl). To a solution of 264 mg of the resultant compound of Example 8D in 10 mL of CH<sub>2</sub>Cl<sub>2</sub> was added 0.14 mL of disopropylethylamine and 175 mg of di-tert butylpyrocarbonate. After stirring for 4 hours, the mixture was diluted with 50 mL of CH<sub>2</sub>Cl<sub>2</sub>, washed with 0.5N of HCl and brine, dried over magnesium sulfate, filtered and concentrated in vacuo to yield 364 mg of the title compound as a white solid which was used without subsequent purification. TLC: R<sub>f</sub> = 0.58, 40% EtOAc/CH<sub>2</sub>Cl<sub>2</sub>.

F. A solution of 334 mg of the resultant compound of Example 8E in 5 mL of ethanol was hydrogenated under 30 psi of hydrogen in the presence of 80 mg of platinum (IV) oxide for 24 hours. The mixture was filtered and concentrated. The residue was purified by a low pressure silica gel column chromatography using 20% EtOAc in CH<sub>2</sub>Cl<sub>2</sub> eluent to yield 268 mg of the title compound. TLC: R<sub>f</sub> = 0.55, 40% EtOAc/CH<sub>2</sub>Cl<sub>2</sub>. (H)-NMR (CDCl<sub>3</sub>) consistent with structure.

G. A solution of 268 mg of the resultant compound of Example 8F in 10 mL of EtOAc was treated with anhydrous HCl gas for 5 min. The reaction mixture was purged with nitrogen then concentrated in vacuo and the

- 76 -

resulting white solid used without subsequent purification for subsequent reaction.

5 H. Compound 195. To a solution of 233 mg of the crude resultant compound of Example 8G in 10 mL of  $\text{CH}_2\text{Cl}_2$  was added 2 mL of saturated aqueous sodium bicarbonate and 149 mg of 4-methyloxybenzene sulfonyl chloride. After 3 hours, the resulting mixture was diluted with  $\text{CH}_2\text{Cl}_2$ , washed with sodium bicarbonate, brine, dried over magnesium sulfate, filtered and  
10 concentrated in vacuo. The residue was purified by low pressure silica gel column chromatography using 0% to 20% EtOAc/ $\text{CH}_2\text{Cl}_2$  to yield 225 mg of the title compound as a white solid. TLC:  $R_f$  = 0.40, 20% EtOAc/ $\text{CH}_2\text{Cl}_2$ ; HPLC:  $R_t$  = 15.65 min.: ( $^1\text{H}$ )NMR ( $\text{CDCl}_3$ ) consistent with  
15 structure.

#### Example 9

##### Synthesis of Compound 196

A. (1S,2 syn)-N-(1-Isobutyl-3-chloro-2-hydroxypropyl)benzyloxycarbonylamine. To a solution of  
20 N-Cbz-leucine chloromethyl ketone (2.0g) in 20 mL of methanol was added, at 0°C, 1.0 g of sodium borohydride and the mixture was stirred at ambient temperature for 24 h. The solution was concentrated under reduced pressure and the residue partitioned between 20 mL of  
25 saturated aqueous  $\text{NH}_4\text{Cl}$  and 500 ml of diethyl ether. The organic fraction was separated, dried over  $\text{MgSO}_4$  and concentrated in vacuo and the residue purified by silica gel chromatography to yield 1.8 g of white solid.

30 B. (1S, 2S)-N-(1-Isobutyl-2,3-epoxypropyl)benzyloxycarbonylamine. To a solution of the resultant compound of Example 9A (300mg) in

- 77 -

absolute ethanol was added 67 mg of powdered KOH. The mixture was stirred for 3 h at ambient temperature, filtered through diatomaceous earth, and concentrated in vacuo. The residue was dissolved in diethyl ether, dried over  $\text{MgSO}_4$ , and concentrated to yield 230 mg of colorless oil, which was used directly for subsequent reaction.

C. (2R,3S)-N<sup>3</sup>-Carbobenzyloxy-N<sup>1</sup>-isobutyl-1,3-diamino-2-hydroxy-5-methylhexane. A 230 mg portion of the resultant compound of Example 9B was suspended in 5 mL of isobutylamine and the mixture stirred overnight at ambient temperature. The mixture was concentrated in vacuo to yield the title product as 179 mg of a white solid, which was used directly for subsequent reaction.

D. Compound V (W = benzyloxycarbonyl, Q = H, D = isobutyl, D' = isobutyl, R' = H, E = 4-methoxyphenyl, (S)-hydroxy). Following the procedure described in Example 8H, a solution of the resultant compound of Example 9C (170mg) in  $\text{CH}_2\text{Cl}_2$  was reacted with 4-methoxybenzenesulfonyl chloride (150 mg) in the presence of aqueous  $\text{NaHCO}_3$ . Workup and silica gel chromatography yielded 90 mg of product as a white solid.

E. Compound V (W = H, Q = H, D = isobutyl, D' = isobutyl, R' = H, E = 4-methoxyphenyl, (syn)-hydroxy). A solution of the resultant compound of Example 9D (90 mg) in ethanol was treated with 50 mg of 10% palladium on carbon and the mixture stirred under an atmosphere of hydrogen. After completion of reaction, the mixture was filtered and concentrated in vacuo to yield 60 mg of the title compound which was used directly for subsequent reaction.

- 78 -

F. Compound 196. Reaction of the resultant compound of Example 9E (60mg) in  $\text{CH}_2\text{Cl}_2$  was reacted with THF-OSu (150 mg) as described earlier yielded, following aqueous workup, drying over  $\text{MgSO}_4$ , filtering, and concentration in vacuo, a residue which was purified by silica gel chromatography using methanol/ $\text{CH}_2\text{Cl}_2$  as eluant to yield the title product as 40 mg of white solid.  $^1\text{H}$ -NMR( $\text{CDCl}_3$ ) consistent with structure.

#### Example 10

##### Synthesis of Compound 203

A. The epoxide prepared in Example 9B (0.430 g, 1.63 mmol) and cyclopentylmethylamine (2.50 g, 25.0 mmol) were stirred at R.T. for 48 h. The solution was diluted with 25 mL of ethanol and concentrated under reduced pressure. The crude material was purified via MPLC (gradient:  $\text{CH}_2\text{Cl}_2$ ; 1% MeOH/ $\text{CH}_2\text{Cl}_2$ ;  $\text{CH}_2\text{Cl}_2$ /MeOH/ $\text{NH}_4\text{OH}$ , 95:5:1) to give 430 mg (73%) of the amine product.  $^1\text{H}$ -NMR( $\text{CDCl}_3$ ) consistent with structure.

B. The amine prepared in Example 10A (0.120 g, 0.331 mmol) was subjected to the procedure described in Example 7B. Purification of the crude material via MPLC (gradient:  $\text{CH}_2\text{Cl}_2$ ; 2%  $\text{Et}_2\text{O}/\text{CH}_2\text{Cl}_2$ ) gave 10 mg of Fraction A (2 syn, 3S isomer) and 70 mg of Fraction B (2R, 3S isomer).  $^1\text{H}$ -NMR( $\text{CDCl}_3$ ) consistent with structures.

C. The Cbz-amine Fraction A prepared in Example 10B (0.010 g, 0.019 mmol) was taken up in 5 mL of absolute ethanol and Pd/C (15 mg) was added. The mixture was stirred under an atmosphere of hydrogen for 24 h. The solution was filtered and concentrated under reduced pressure to give the amine product in quantitative



- 79 -

yield. This material was used without further purification.

- 5 D. The amine prepared in Example 10C (0.010 g, 0.025 mmol) was used in the procedure described in Example 1D. Purification of the crude material via MPLC (gradient:  $\text{CH}_2\text{Cl}_2$ ; 1%  $\text{MeOH}/\text{CH}_2\text{Cl}_2$ ) gave 2.8 mg (29%) of compound 203.

#### Example 11

##### Synthesis of Compound 212

- 10 A. A 1.12 portion of anhydrous DMF is cooled to 0 °C and treated dropwise with 2.06 g of sulfuryl chloride. The resulting suspension is stirred for 30 min., then treated with 1.50 g of benzyl phenyl ether. The mixture is heated at 90 °C for 3h, then cooled,  
15 extracted with brine and methylene chloride, and dried over  $\text{MgSO}_4$ . Chromatography on silica gel (i- $\text{PrOH}/\text{hexanes}$ ) yields 4-benzyloxybenzenesulfonyl chloride.
- 20 B. The amine prepared in Example 7A (0.150 g, 0.398 mmol) and 4-(phenylmethoxy)benzenesulfonyl chloride (0.170 g, 0.601 mmol) were utilized in the procedure described in Example 7B to give after MPLC purification (gradient:  $\text{CH}_2\text{Cl}_2$ ; 5%  $\text{Et}_2\text{O}/\text{CH}_2\text{Cl}_2$ ; 10%  $\text{Et}_2\text{O}/\text{CH}_2\text{Cl}_2$ ) 120 mg (48%) of Cbz-amine.  $^1\text{H}$ -NMR( $\text{CDCl}_3$ ) consistent with  
25 structure.
- C. The Cbz-amine prepared in Example 11B (0.120 g, 0.214 mmol) was subjected to the procedure described in Example 10C to give 50 mg (59%) of the crude amine which was used without further purification.
- 30 D. The amine prepared in Example 11C (0.050 g, 0.125 mmol) was used in the procedure described in Example

- 80 -

1D. Purification of the crude material via MPLC (gradient: CH<sub>2</sub>Cl<sub>2</sub>; 1% MeOH/CH<sub>2</sub>Cl<sub>2</sub>; 2% MeOH/CH<sub>2</sub>Cl<sub>2</sub>) gave compound 212. [<sup>1</sup>H]-NMR(CDCl<sub>3</sub>) consistent with structure.

5

Example 12Synthesis of Compound 213

- A. The epoxide described in Example 9B (0.53 g, 2.01 mmol) was reacted with 2-phenylethylamine (5.0 mL, 40 mmol) as described in Example 9C to give 640 mg (83% yield) of the amine after MPLC purification (gradient: CH<sub>2</sub>Cl<sub>2</sub>; 1% then 5% then 10% MeOH/CH<sub>2</sub>Cl<sub>2</sub>).  
10
- B. The amine prepared in Example 12A (0.150 g, 0.39 mmol) was subjected to the procedure described in Example 11B to give after MPLC purification (gradient: CH<sub>2</sub>Cl<sub>2</sub>; 5% then 10% methanol/CH<sub>2</sub>Cl<sub>2</sub>) 110 mg (45%) of the Cbz-amine. [<sup>1</sup>H]-NMR(CDCl<sub>3</sub>) consistent with structure.  
15
- C. The Cbz-amine prepared in Example 12B (0.110 g, 0.177 mmol) was treated as described in Example 10C to give 40 mg (56%) of the amine which was used without further purification. [<sup>1</sup>H]-NMR(CDCl<sub>3</sub>) consistent with structure.  
20
- D. The amine prepared in Example 12C (0.040 g, 0.098 mmol) was treated as described in Example 1D. Purification of the crude material via MPLC (gradient: CH<sub>2</sub>Cl<sub>2</sub>; 1% MeOH/CH<sub>2</sub>Cl<sub>2</sub>; 2% MeOH/CH<sub>2</sub>Cl<sub>2</sub>) gave compound 213. [<sup>1</sup>H]-NMR(CDCl<sub>3</sub>) consistent with structure.  
25

Example 13Synthesis of Compound 223

5 A. The procedure described in Example 1A was performed using 30 mg of epoxide VI (W=Boc, Q=H) and 2-pyrrolidinylethylamine (0.200 g, 1.75 mmol) to give 25 mg (58%) of Boc-amine after chromatography (CHCl<sub>3</sub>/MeOH, 2:1). [<sup>1</sup>H]-NMR(CDCl<sub>3</sub>) consistent with structure.

10 B. The procedure described in Example 7B was performed using the amine prepared in Example 13A (0.040 g, 0.106 mmol) and substituting potassium bicarbonate for sodium bicarbonate (in both solid and aqueous solution forms) to give 30 mg (52%) of BOC-amine sulfonamide after chromatography (CHCl<sub>3</sub>/MeOH, 5:1). [<sup>1</sup>H]-NMR(CDCl<sub>3</sub>) consistent with structure.

15 C. The BOC-amine sulfonamide prepared in Example 13B (0.008 g, 0.015 mmol) was dissolved in acetonitrile and treated with 2N HCl. The solvents were removed and the material dried over MgSO<sub>4</sub>. This crude material was subjected to the procedure described in Example 1D to give 5 mg (59%) of compound 223 after chromatography (CHCl<sub>3</sub>/MeOH, 10:1). [<sup>1</sup>H]-NMR(CDCl<sub>3</sub>) consistent with structure.

20

#### Example 14

##### Synthesis of Compound 224

25 A. Cyclopentylacetic acid (5.2 g, 41 mmol) and thionyl chloride (10 mL) were combined then DMF (0.2 mL) was added and the solution was stirred 1.5 h at R.T. Dichloromethane (10 mL) was added and the solution cooled in an ice bath whereupon 25% aqueous ammonia (30 mL) was added and the mixture stirred for 0.5 h. The solution was extracted with CH<sub>2</sub>Cl<sub>2</sub> (3X) and the combined extracts washed with 1N HCl, dried over MgSO<sub>4</sub>, and concentrated to give cyclopentylacetamide

30

- 82 -

(2.699 g, 52%).  $^1\text{H}$ -NMR( $\text{CDCl}_3$ ) consistent with structure.

5 B. Cyclopentylacetamide (2.70 g, 21.0 mmol) was dissolved in 100 mL of  $\text{Et}_2\text{O}$  whereupon lithium aluminum hydride (2.2 g, 58 mmol) was added and the mixture was heated to 60 °C for 4 h. Following standard work-up conditions the crude material was distilled to give 2-cyclopentylethylamine (750 mg, 32%); b.p.: 78 °C at 40 mm Hg.  $^1\text{H}$ -NMR( $\text{CDCl}_3$ ) consistent with structure.

10 C. The procedure described in Example 1A was performed using 14 mg of epoxide VI ( $\text{W}=\text{Boc}$ ,  $\text{Q}=\text{H}$ ) and 2-cyclopentylethylamine (0.045 g, 0.40 mmol) to give 12 mg (64%) of Boc-amine after chromatography ( $\text{CHCl}_3/\text{MeOH}/\text{NH}_4\text{OH}$ ; 90:10:1).  $^1\text{H}$ -NMR( $\text{CDCl}_3$ ) consistent  
15 with structure.

D. The BOC-amine prepared in Example 14C (0.188 g, 0.5 mmol) was dissolved in  $\text{CH}_2\text{Cl}_2$  and TEA (0.09 mL) and 4-methoxybenzenesulfonyl chloride (0.124 g, 0.6 mmol) were added. After stirring for 2 h the mixture was  
20 washed with saturated aq.  $\text{NaHCO}_3$  (3X), 10% aq.  $\text{HCl}$  (3X), and dried over  $\text{MgSO}_4$  to give 240 mg (88%) of BOC-amine sulfonamide after chromatography (hexanes/ $\text{EtOAc}$ ; 8:2).  $^1\text{H}$ -NMR( $\text{CDCl}_3$ ) consistent with structure.

E. The BOC-amine sulfonamide prepared in Example 14D  
25 (0.235 g, 0.43 mmol) was taken up in  $\text{EtOAc}$  and treated with  $\text{HCl}$  to give 214 mg (quantitative) of crude amine- $\text{HCl}$  salt which was used without further purification.

F. The crude amine- $\text{HCl}$  salt prepared in Example 14E (0.211 g, 0.43 mmol) was treated as described in

- 83 -

Example 1D to give 178 mg (74%) of compound 224. <sup>1</sup>H-NMR(CDCl<sub>3</sub>) consistent with structure.

Example 15

Synthesis of Compound 225

- 5     A.    4-Piperidinecarboxamide (10.2 g, 78.0 mmol) in DMSO (10 mL) was treated with benzyl bromide (20.0 mL, 168 mmol). The mixture was diluted with EtOAc and washed with 1N aq. HCl, 5N aq. NaOH, dried over MgSO<sub>4</sub>, and concentrated to give 5.90 g (35%) of N-benzyl-4-
- 10    piperidinecarboxamide. <sup>1</sup>H-NMR(CDCl<sub>3</sub>) consistent with structure.
- 15    B.    The N-benzyl-4-piperidinecarboxamide (2.19 g, 10 mmol) was treated as described in Example 14B to give 2.01 g (98%) of the amine. <sup>1</sup>H-NMR(CDCl<sub>3</sub>) consistent with structure.
- 20    C.    The procedure described in Example 1A was performed using 100 mg of epoxide VI (W=Boc, Q=H) and the amine prepared in Example 15B (0.100 g, 0.78 mmol) to give 25 mg (14%) of Boc-amine after chromatography (hexanes:EtOAc; 10:1). <sup>1</sup>H-NMR(CDCl<sub>3</sub>) consistent with structure.
- 25    D.    The procedure described in Example 14D was performed using the BOC-amine prepared in Example 15C (0.106 g, 0.23 mmol) to give 92 mg (65%) of the sulfonamide. <sup>1</sup>H-NMR(CDCl<sub>3</sub>) consistent with structure.
- E.    The sulfonamide prepared in Example 15D (0.065 g, 0.10 mmol) was treated as described in Example 14E to give 64 mg of crude amine-HCl salt. <sup>1</sup>H-NMR(CDCl<sub>3</sub>) consistent with structure.

- 84 -

F. The crude amine-HCl salt prepared in Example 15E (0.055 g, 0.092 mmol) was treated as described in Example 1D to give 51 mg (85%) of N-benzylpiperidine.  $^1\text{H}$ -NMR( $\text{CDCl}_3$ ) consistent with structure.

5 G. The N-benzylpiperidine (0.055 g, 0.092 mmol), prepared in Example 15F was treated as described in Example 10C to give a crude product which upon purification by chromatography ( $\text{CH}_2\text{Cl}_2/\text{MeOH}/\text{NH}_4\text{OH}$ ; 90:10:1) gave compound 225.  $^1\text{H}$ -NMR( $\text{CDCl}_3$ ) consistent  
10 with structure.

#### Example 16

##### Synthesis of Compound 228

A. 4-Hydroxypiperidine (10.8 g, 107 mmol) and  $\text{Et}_3\text{N}$  (17 mL, 123 mmol) in 80 mL of  $\text{CH}_2\text{Cl}_2$  was cooled in an  
15 ice bath whereupon benzyl chloroformate (16.3 mL, 114 mmol) was added. After stirring 1.5 h at R.T. the mixture was subjected to standard work-up conditions to give 13.9 g (55%) of N-Cbz-piperidine after chromatography ( $\text{CHCl}_3/\text{MeOH}$ ; 10:1).  $^1\text{H}$ -NMR( $\text{CDCl}_3$ )  
20 consistent with structure.

B. A solution of DEAD (1.61 mL, 10.2 mmol) in 20 mL of THF was added to a solution of triphenylphosphine (2.69 g, 10.3 mmol), the N-Cbz-piperidine prepared in Example 16A (2.36 g, 10.0 mmol) and phthalimide (1.50 g, 10.2 mmol) in 80 mL of THF. After stirring 10.5 h at  
25 R.T. the mixture was quenched with water, extracted with EtOAc (3X) and the combined extracts washed with brine solution, dried over  $\text{Na}_2\text{SO}_4$  and concentrated to give the crude product. Purification of this material  
30 by chromatography (hexanes/EtOAc; 2:1) gave 1.81 g, (50%) of 1-benzoyloxycarbonyl-4-phthalimidylpiperidine.  $^1\text{H}$ -NMR( $\text{CDCl}_3$ ) consistent with structure.

- 85 -

- C. 1-Benzyloxycarbonyl-4-phthalimidylpiperidine (1.50 g, 4.27 mmol), prepared in Example 16B, was taken up in 20 mL of ethanol. To this solution was added hydrazine monohydrate (35 mL, 700 mmol) and this mixture was  
5 heated at 100 °C for 3 h. Brine solution (40 mL) and 10% aq.  $K_2CO_3$  (60 mL) were added and the mixture was extracted with 5% MeOH/ $CHCl_3$  (3X). The combined extracts were washed with 2N aq. HCl, 2N aq. NaOH, brine solution, dried over  $Na_2SO_4$  and concentrated to  
10 give 0.847 g, (85%) of 4-amino-1-benzyloxycarbonylpiperidine.  $[^1H]$ -NMR( $CDCl_3$ ) consistent with structure.
- D. The procedure described in Example 1A was performed using 132 mg of epoxide VI (W=Boc, Q=H) and  
15 4-amino-1-benzyloxycarbonylpiperidine (0.353 g, 1.51 mmol), prepared in Example 16C, to give 168 mg (67%) of Boc-amine after chromatography ( $CHCl_3$ /MeOH/ $NH_4OH$ ; 95:5:1).  $[^1H]$ -NMR( $CDCl_3$ ) consistent with structure.
- E. The procedure described in Example 14D was  
20 performed using the BOC-amine prepared in Example 16D (0.059 g, 0.320 mmol) to give the crude product which after chromatography ( $CHCl_3$ /MeOH/ $NH_4OH$ ; 95:5:1) gave 141 mg (66%) of the sulfonamide.  $[^1H]$ -NMR( $CDCl_3$ ) consistent with structure.
- F. The sulfonamide prepared in Example 16E (0.050 g, 1.57 mmol) was treated as described in Example 14E to  
25 give 971 mg of crude amine-HCl salt.  $[^1H]$ -NMR( $CDCl_3$ ) consistent with structure.
- G. The crude amine-HCl salt prepared in Example 16F  
30 (0.118 g, 0.207 mmol) was treated as described in

- 86 -

Example 1D to give 147 mg of crude carbamate.  $^1\text{H}$ -NMR( $\text{CDCl}_3$ ) consistent with structure.

H. The crude carbamate prepared in Example 16G (0.116 g, 0.163 mmol) was treated as described in Example 10C. Purification of the crude material by chromatography ( $\text{CHCl}_3/\text{MeOH}/\text{NH}_4\text{OH}$ ; 95:5:1) gave 28 mg (31%) of the compound 228.  $^1\text{H}$ -NMR( $\text{CDCl}_3$ ) consistent with structure.

#### Example 17

##### Synthesis of Compound 233

A. The procedure described in Example 1A was performed using epoxide VI (W=Boc, Q=H) (0.030 g, 0.11 mmol) and 2,2-dimethyl-3-hydroxypropylamine (0.24 g, 0.23 mmol) to give 42 mg (quantitative) of Boc-amine after chromatography ( $\text{CHCl}_3/\text{MeOH}/\text{NH}_4\text{OH}$ ; 100:10:1).  $^1\text{H}$ -NMR( $\text{CDCl}_3$ ) consistent with structure.

B. The procedure described in Example 14D was performed using the BOC-amine prepared in Example 17A (0.030 g, 0.082 mmol) to give the crude product which after chromatography ( $\text{CHCl}_3/\text{MeOH}/\text{NH}_4\text{OH}$ ; 150:10:1) gave 42.8 mg (95%) of the sulfonamide.  $^1\text{H}$ -NMR( $\text{CDCl}_3$ ) consistent with structure.

C. The sulfonamide prepared in Example 17B (0.030 g, 0.056 mmol) was treated as described in Example 14E to give 29.3 mg of crude amine-HCl salt.  $^1\text{H}$ -NMR( $\text{CDCl}_3$ ) consistent with structure.

D. The crude amine-HCl salt prepared in Example 17C (0.029 g, 0.061 mmol) was treated as described in Example 1D to give 21.1 mg (69%) of compound 233.  $^1\text{H}$ -NMR( $\text{CDCl}_3$ ) consistent with structure.



- 87 -

Example 18Synthesis of Compound 234

5 A. The procedure described in Example 1A was performed using epoxide VI (W=Boc, Q=H) (0.200 g, 0.76 mmol) and 2-aminothiazole (1.2 g, 12.0 mmol) to give 150 mg (54%) of Boc-amine after chromatography (CHCl<sub>3</sub>/MeOH; 10:1). [<sup>1</sup>H]-NMR(CDCl<sub>3</sub>) consistent with structure.

10 B. The procedure described in Example 14D was performed using the BOC-amine prepared in Example 18A (0.004 g, 0.011 mmol) to give the crude product which after chromatography (CHCl<sub>3</sub>/EtOAc; 3:4) gave the sulfonamide. [<sup>1</sup>H]-NMR(CDCl<sub>3</sub>) consistent with structure.

15 C. The sulfonamide prepared in Example 18B (0.010 g, 0.019 mmol) was treated as described in Example 14E to give the crude amine-HCl salt. This material was subjected to the procedure described in Example 1D to give 6 mg (58% for two steps) of compound 234 after chromatography (CHCl<sub>3</sub>/MeOH; 10:1). ([<sup>1</sup>H]-NMR(CDCl<sub>3</sub>) consistent with structure.

20

Example 19Synthesis of Compound 235

25 A. 4-Methoxybenzenesulfonyl chloride (0.035 g, 0.169 mmol) was dissolved in pyridine and 4-amino-1,2,4-triazole (0.17 g, 0.202 mmol) was added. After 4 days at R.T. the mixture was subjected to standard work-up conditions to give 33 mg of sulfonamide.

30 B. The sulfonamide prepared in Example 19A (0.356 g, 0.140 mmol) was treated with KOH (0.078 g, 0.139 mmol) to give the corresponding potassium salt.

- 88 -

C. The epoxide VI (W=Boc, Q=H) (0.049 g, 0.168 mmol) and the potassium salt of the amine prepared in Example 19B (0.044 g, 0.169 mmol) were reacted in DMSO at 80 °C for 2 days to give 9 mg (10%) of Boc-amine. [<sup>1</sup>H]-  
5 NMR(CDCl<sub>3</sub>) consistent with structure.

D. The sulfonamide prepared in Example 19C (0.143 g, 0.276 mmol) was treated as described in Example 14E to give the crude amine-HCl salt. This material was subjected to the procedure described in Example 1D to  
10 give 97 mg (66% for two steps) of compound 235 after recrystallization (CH<sub>2</sub>Cl<sub>2</sub>/MeOH). ([<sup>1</sup>H]-NMR(CDCl<sub>3</sub>) consistent with structure.

#### Example 20

##### Synthesis of Compounds 167 and 168

15 A. Compound 167. A solution of 102 mg of N-((2 syn, 3S)-2-hydroxy-4-phenyl-3-((S)-tetrahydrofuran-3-yloxy-carbonylamino)-butyl)-N-isobutylbenzenesulfonamide in 4:1 CH<sub>2</sub>Cl<sub>2</sub>/saturated aqueous NaHCO<sub>3</sub> was treated sequentially, at ambient temperature under an  
20 atmosphere of nitrogen, with 65 mg of p-nitrobenzenesulfonyl chloride and 51 mg of sodium bicarbonate. The mixture was stirred for 14 h, diluted with CH<sub>2</sub>Cl<sub>2</sub>, washed with saturated NaCl, then dried over MgSO<sub>4</sub>, filtered, and concentrated in vacuo. The residue  
25 was purified by low pressure silica gel chromatography using 20% diethyl ether/CH<sub>2</sub>Cl<sub>2</sub> as eluent to provide 124 mg of the title product as a white solid. TLC: R<sub>f</sub> = 0.36, 20% diethyl ether/CH<sub>2</sub>Cl<sub>2</sub>. HPLC: R<sub>t</sub> = 15.15 min. [<sup>1</sup>H]-NMR (CDCl<sub>3</sub>) consistent with structure.

30 B. Compound 168. A solution of 124 mg of compound 167 in ethyl acetate was treated, at ambient

- 89 -

temperature, with 13 mg of 10% palladium on carbon. The mixture was stirred for 14 h under an atmosphere of hydrogen, filtered through a pad of Celite filter agent, and concentrated in vacuo. The residue was  
5 subjected to preparative HPLC to yield 82 mg of the title product as a white solid. TLC:  $R_f = 0.10$ , 20% ether/ $\text{CH}_2\text{Cl}_2$ . HPLC:  $R_t = 13.16$  min. ( $^1\text{H}$ )-NMR ( $\text{CDCl}_3$ ) consistent with structure.

#### Example 21

10 We measured the inhibition constants of the compounds listed in Table II against HIV-1 protease using the above-cited method of Pennington et al.

We also measured the anti-viral potency of the compounds in CCRM-CEM cells by the above-cited  
15 method of Meek et al. The results are shown in Table III. In the Tables below,  $K_i$  and  $\text{IC}_{50}$  values are expressed in nM. The designation "ND" is used where a given compound was not tested.

20 In Table III, the following classifications have been employed:

- A: inhibits HIV replication at concentration of 100 nM or less.
- B: inhibits HIV replication at concentration of between 101 and 1,000 nM.
- 25 C: inhibits HIV replication at a concentration of between 1,001 and 10,000 nM.
- D: inhibits HIV replication at a concentration of between 10,001 and 40,000 nM.

- 90 -

TABLE II

	<u>Compound</u>	<u>K<sub>i</sub></u>	<u>Compound</u>	<u>K<sub>i</sub></u>	<u>Compound</u>	<u>K<sub>i</sub></u>
	35	4.0	168	0.6	229	11.0
	37	0.1	169	<0.1	230	9,300
5	48	1.4	170	0.2	231	1,400
	51	ND	171	0.2	232	20.0
	52	0.4	172	21	233	4.0
	53	27	173	0.6	234	10,000
	54	22	174	10	235	1,700
10	60	4.0	175	0.1	236	700.0
	66	0.4	176	<0.1	237	ND
	69	42	180	<0.1		
	86	5.0	181	0.3		
	88	1.4	182	0.2		
15	91	2.5	183	0.1		
	93	0.8	195	0.2		
	94	1.7	196	ND		
	95	1.3	197	<.1		
	99	0.24	198	<.1		
20	100	0.16	199	10.0		
	101	250	200	2.4		
	112	4.0	201	7.5		
	113	3.0	202	.1		
	116	<0.1	203	60.0		
25	123	10	204	4.0		
	124	1.1	205	5.0		
	125	0.3	206	.50		
	132	6.0	207	33.0		
	133	24	208	5.5		
30	134	8.4	209	.8		
	135	2.7	210	6.0		
	136	18	211	<0.1		
	137	26	212	1.0		
	138	1.4	213	5.0		
35	140	<0.1	214	110.0		
	144	8.0	215	11.0		
	145	1.4	216	<0.1		
	148	0.2	217	0.3		
	149	1.7	218	5.0		
40	150	6.0	219	1.0		
	151	0.8	220	0.2		
	152	2.5	221	1.0		
	157	0.7	222	0.5		
	158	<0.1	223	3,800		
45	159	0.2	224	1.0		
	160	1.0	225	1,500		
	161	20	226	3,000		
	165	0.4	227	9.0		
	167	0.45	228	4,550		

- 91 -

TABLE III

	<u>Compound</u>	<u>IC<sub>90</sub></u>	<u>Compound</u>	<u>IC<sub>90</sub></u>
5	35	B	169	A
	37	B	170	B
	48	B	171	A
	51	C	172	ND
	52	B	173	A
10	53	ND	174	ND
	60	C	175	A
	66	B	176	ND
	69	ND	180	ND
	86	B	181	ND
15	88	B	182	B
	91	B	183	B
	93	B	195	A
	94	B	196	ND
	95	C	197	ND
20	99	B	198	ND
	100	A	199	ND
	101	ND	200	ND
	112	B	201	ND
	113	B	202	ND
25	116	A	203	ND
	123	ND	204	ND
	124	D	205	ND
	125	B	206	B
	132	ND	207	ND
30	133	ND	208	ND
	134	ND	209	B
	135	C	210	ND
	136	ND	211	A
	137	ND	212	B
35	138	B	213	ND
	140	A	214	ND
	144	B	215	ND
	145	B	216	A
	148	A	217	A
40	149	B	218	ND
	150	B	219	A
	151	C	220	A
	152	ND	221	B
	157	B	222	ND
45	158	A	223	ND
	159	B	224	ND
	160	A	225	ND
	161	ND	226	ND
	165	B	227	ND
	167	B	228	ND
	168	A	229	ND

	<u>Compound</u>	<u>IC<sub>90</sub></u>
	230	ND
	231	ND
	232	ND
5	233	ND
	234	ND
	235	ND
	236	ND
	237	ND


10           As demonstrated in Tables II and III, all of the compounds tested displayed inhibitory and anti-viral activity. Moreover, several of these compounds exhibited activity levels which are among the highest levels known to date for HIV protease inhibitors.

15           While we have described a number of embodiments of this invention, it is apparent that our basic constructions may be altered to provide other embodiments which utilize the products and methods of this invention. Therefore, it will be appreciated that  
20           the scope of this invention is to be defined by the appended claims, rather than by the specific embodiments which have been presented by way of example.

CLAIMS

We claim:

1. A compound of formula I:



wherein:

each  $\text{R}^1$  is independently selected from the group consisting of  $-\text{C}(\text{O})-$ ,  $-\text{S}(\text{O})_2-$ ,  $-\text{C}(\text{O})-\text{C}(\text{O})-$ ,  $-\text{O}-\text{C}(\text{O})-$ ,  $-\text{O}-\text{S}(\text{O})_2-$ ,  $-\text{NR}^2-\text{S}(\text{O})_2-$ ,  $-\text{NR}^2-\text{C}(\text{O})-$  and  $-\text{NR}^2-\text{C}(\text{O})-\text{C}(\text{O})-$ ;

each Het is independently selected from the group consisting of  $\text{C}_3\text{-C}_7$  carbocycle;  $\text{C}_6\text{-C}_{10}$  aryl; phenyl fused with heterocycle; and heterocycle; wherein any member of said Het may be optionally substituted with one or more substituents selected from the group consisting of oxo,  $-\text{OR}^2$ ,  $-\text{R}^2$ ,  $-\text{N}(\text{R}^2)(\text{R}^2)$ ,  $-\text{NHOH}$ ,  $-\text{R}^2\text{-OH}$ ,  $-\text{CN}$ ,  $-\text{CO}_2\text{R}^2$ ,  $-\text{C}(\text{O})-\text{N}(\text{R}^2)(\text{R}^2)$ ,  $-\text{S}(\text{O})_2-\text{N}(\text{R}^2)(\text{R}^2)$ ,  $-\text{N}(\text{R}^2)-\text{C}(\text{O})-\text{R}^2$ ,  $-\text{C}(\text{O})-\text{R}^2$ ,  $-\text{S}(\text{O})_n-\text{R}^2$ ,  $-\text{OCF}_3$ ,  $-\text{S}(\text{O})_n-\text{R}^6$ ,  $-\text{N}(\text{R}^2)-\text{S}(\text{O})_2(\text{R}^2)$ , halo,  $-\text{CF}_3$ ,  $-\text{NO}_2$ ,  $-\text{R}^6$  and  $-\text{O}-\text{R}^6$ ;

each  $\text{R}^2$  is independently selected from the group consisting of H and  $\text{C}_1\text{-C}_3$  alkyl optionally substituted with  $\text{R}^6$ ;

each  $\text{R}^3$  is independently selected from the group consisting of H, Het,  $\text{C}_1\text{-C}_6$  alkyl and  $\text{C}_2\text{-C}_6$  alkenyl wherein any member of said  $\text{R}^3$ , except H, may be optionally substituted with one or more substituents selected from the group consisting of  $-\text{OR}^2$ ,  $-\text{C}(\text{O})-\text{NH}-\text{R}^2$ ,  $-\text{S}(\text{O})_n-\text{N}(\text{R}^2)(\text{R}^2)$ , Het,  $-\text{CN}$ ,  $-\text{SR}^2$ ,  $-\text{CO}_2\text{R}^2$ ,  $\text{NR}^2-\text{C}(\text{O})-\text{R}^2$ ;

each n is independently 1 or 2;

each D and  $\text{D}'$  is independently selected from the group consisting of  $\text{R}^6$ ;  $\text{C}_1\text{-C}_3$  alkyl, which may be optionally substituted with one or more groups selected from  $-\text{OR}^2$ ,  $-\text{R}^3$ ,  $-\text{O}-\text{R}^6$ ,  $-\text{S}-\text{R}^6$  and  $\text{R}^6$ ;  $\text{C}_2\text{-C}_6$  alkenyl, which

- 94 -

may be optionally substituted with one or more groups selected from the group consisting of  $-OR^2$ ,  $-R^3$ ,  $-O-R^6$  and  $R^6$ ;  $C_3-C_6$  carbocycle, which may be optionally substituted with or fused with  $R^6$ ; and  $C_1-C_6$  cycloalkenyl, which may be optionally substituted with or fused with  $R^6$ ;

each E is independently selected from the group consisting of Het;  $-O-Het$ ; Het-Het;  $-O-R^3$ ;  $-NR^2R^3$ ;  $C_1-C_6$  alkyl, which may be optionally substituted with one or more groups selected from the group consisting of  $R^4$  and Het; and  $C_2-C_6$  alkenyl, which may be optionally substituted with one or more groups selected from the group consisting of  $R^4$  and Het;

each  $R^4$  is independently selected from the group consisting of  $-OR^2$ ,  $-C(O)-NHR^2$ ,  $-S(O)_2-NHR^2$ , halo,  $-NR^2-C(O)-R^2$  and  $-CN$ ;

each  $R^5$  is independently selected from the group consisting of H and  $C_1-C_4$  alkyl optionally substituted with aryl; and

each  $R^6$  is independently selected from the group consisting of aryl, carbocycle and heterocycle, wherein said carbocycle or heterocycle may be optionally substituted with one or more groups selected from the group consisting of oxo,  $-OR^5$ ,  $-R^5$ ,  $-N(R^5)(R^5)$ ,  $-N(R^5)-C(O)-R^5$ ,  $-R^5-OH$ ,  $-CN$ ,  $-CO_2R^5$ ,  $-C(O)-N(R^5)(R^5)$ , halo and  $-CF_3$ .

2. The compound according to claim 1, wherein THF is tetrahydrofuran-3-yl.

3. The compound according to claim 1, wherein  $R^1$  is  $-O-C(O)-$ .

4. The compound according to claim 1, wherein D is selected from the group consisting of



- 95 -

benzyl, isobutyl, cyclopentylmethyl, cyclohexylmethyl and phenylmethyl.

5. The compound according to claim 1, wherein:

each D' is independently selected from the group consisting of C<sub>1</sub>-C<sub>4</sub> alkyl optionally substituted with R<sup>6</sup>;

each R<sup>6</sup> is independently selected from the group consisting of aryl, 3-6 membered carbocycle and 5-6 membered heterocycle, wherein said heterocycle or carbocycle may be optionally substituted with one or more groups selected from the group consisting of oxo, -OR<sup>5</sup>, -R<sup>5</sup>, -N(R<sup>5</sup>)(R<sup>5</sup>), -N(R<sup>5</sup>)-C(O)-R<sup>5</sup>, -R<sup>5</sup>-OH, -CN, -CO<sub>2</sub>R<sup>5</sup>, -C(O)-N(R<sup>5</sup>)(R<sup>5</sup>), halo and -CF<sub>3</sub>; and

each R<sup>5</sup> is independently selected from the group consisting of H and C<sub>1</sub>-C<sub>4</sub> alkyl optionally substituted with aryl.

6. The compound according to claim 5, wherein D' is selected from the group consisting of C<sub>1</sub>-C<sub>4</sub> alkyl optionally substituted with one 3-6 membered carbocycle or one 5-6 membered heterocycle.

7. The compound according to claim 6, wherein D' is selected from the group consisting of isobutyl, cyclopentylmethyl and cyclohexylmethyl.

8. The compound according to claim 1, wherein:

each E is independently phenyl optionally substituted with one or more substituents selected from the group consisting of -OR<sup>2</sup>, -R<sup>2</sup>, -N(R<sup>2</sup>)(R<sup>2</sup>), -N(R<sup>2</sup>)-C(O)-R<sup>2</sup>, -R<sup>2</sup>-OH, -CN, -CO<sub>2</sub>R<sup>2</sup>, -C(O)-N(R<sup>2</sup>)(R<sup>2</sup>), halo, and

- 96 -

-CF<sub>3</sub>; or phenyl fused with a 5-7 membered heterocycle or carbocycle;

each R<sup>2</sup> is independently selected from the group consisting of H and C<sub>1</sub>-C<sub>3</sub> alkyl optionally substituted with R<sup>6</sup>;

each R<sup>6</sup> is independently selected from the group consisting of aryl, 3-6 membered carbocycle and 5-6 membered heterocycle, wherein said carbocycle or heterocycle may be optionally substituted with one or more groups selected from the group consisting of oxo, -OR<sup>5</sup>, -R<sup>5</sup>, -N(R<sup>5</sup>)(R<sup>5</sup>), -N(R<sup>5</sup>)-C(O)-R<sup>5</sup>, -R<sup>5</sup>-OH, -CN, -CO<sub>2</sub>R<sup>5</sup>, -C(O)-N(R<sup>5</sup>)(R<sup>5</sup>), halo and -CF<sub>3</sub>; and

each R<sup>5</sup> is independently selected from the group consisting of H and C<sub>1</sub>-C<sub>4</sub> alkyl optionally substituted with aryl.

9. The compound according to claim 8, wherein E is phenyl substituted with one or more substituents selected from the group consisting of -OH, -OCH<sub>3</sub>, -NH<sub>2</sub>, -NHCOCH<sub>3</sub>, -SH, and -CH<sub>3</sub>; or phenyl fused with 5-6 membered heterocycle or carbocycle.

10. The compound according to claim 9, wherein E is phenyl substituted with -NH<sub>2</sub> at the meta- or para-position.

11. The compound according claim 1, wherein the configuration of the chiral carbon of THF is (S).

12. A compound selected from the group consisting of: compound 35; compound 37; compound 48; compounds 52; compound 60; compound 66; compounds 86; compound 88; compound 91; compound 93; compound 94; compound 95; compound 99; compound 100; compound 112; compound 113; compound 116; compound 124; compound 125;

- 97 -

compound 132; compound 134; compound 135; compound 138; compound 140; compound 144; compound 145; compound 148; compound 149; compound 150; compound 151; compound 152; compound 157; compound 158; compound 159; compound 160; compound 165; compound 167; compound 168; compound 169; compound 170; compound 171; compound 173; compound 175; compound 176; compound 180; compound 181; compound 182; compound 183; compound 195; compound 196; compound 197; compound 198; compound 200; compound 201; compound 202; compound 203; compound 204; compound 205; compound 206; compound 208; compound 209; compound 210; compound 211; compound 212; compound 213; compound 216; compound 217; compound 218; compound 219; compound 220; compound 221; compound 222; compound 224; compound 227; and compound 233; wherein each compound has the formula shown in Table I.

13. The compound according to claim 12 selected from the group consisting of: compound 48; compound 100; compound 116, compound 140; compound 148; compound 158; compound 160; compound 168; compound 169; compound 171; compound 173; compound 175; compound 176; compound 180; compound 181; compound 195; compound 197; compound 198; compound 202; compound 211; compound 216; compound 217; compound 219; and compound 220, wherein each compound has the formula shown in Table I.

14. A compound having the formula of compound 140 shown in Table I.

15. A compound having the formula of compound 168 shown in Table I.

- 98 -

16. A compound having the formula of compound 169 shown in Table I.

17. A compound having the formula of compound 171 shown in Table I.

18. A compound having the formula of compound 175 shown in Table I.

19. A compound having the formula of compound 216 shown in Table I.

20. A compound having the formula of compound 217 shown in Table I.

21. A pharmaceutical composition comprising a pharmaceutically effective amount of a compound according to claim 1 and a pharmaceutically acceptable carrier, adjuvant or vehicle.

22. The pharmaceutical composition according to claim 21, wherein said pharmaceutical composition is orally administrable.

23. The pharmaceutical composition according to claim 21, further comprising one or more additional agents selected from the group consisting of other anti-viral agents and immunostimulators.

24. The pharmaceutical composition according to claim 23, wherein said other anti-viral agent or agents are protease inhibitors or reverse transcriptase inhibitors.

- 99 -

25. The pharmaceutical composition according to claim 24, wherein said protease inhibitor or inhibitors are HIV protease inhibitors.

26. The pharmaceutical composition according to claim 25, wherein said HIV protease inhibitor or inhibitors are selected from the group consisting of saquinavir (Ro 31-8959), MK 639, APT 538 (A80538), AG 1343, XM 412, XM 450 and BMS 186,318.

27. The pharmaceutical composition according to claim 24, wherein said reverse transcriptase inhibitor or inhibitors are nucleoside analogs.

28. The pharmaceutical composition according to claim 27, wherein said nucleoside analog or analogs are selected from the group consisting of zidovudine (AZT), dideoxycytidine (ddC), didanosine (ddI), stavudine (d4T), 3TC, 935U83, 1592U89 and 524W91.

29. The pharmaceutical composition according to claim 24, wherein said reverse transcriptase inhibitor or inhibitors are non-nucleoside analogs.

30. The pharmaceutical composition according to claim 29, wherein said non-nucleoside reverse transcriptase inhibitor or inhibitors are delavirdine (U90) or nevirapine.

31. A method for inhibiting aspartyl protease activity comprising the step of contacting an aspartyl protease with the compound according to claim 1.

- 100 -

32. A method for reversibly binding an aspartyl protease comprising the step of contacting the aspartyl protease with the compound according to claim 1, said compound being covalently bound to a solid matrix.

33. A method for preventing HIV infection in a mammal comprising the step of administering to said mammal a pharmaceutically effective amount of a pharmaceutical composition according to either claim 21 or 22.

34. A method for preventing HIV infection in a mammal comprising the step of administering to said mammal a pharmaceutically effective amount of a pharmaceutical composition according to claim 23.

35. A method for treating HIV infection in a mammal comprising the step of administering to said mammal a pharmaceutically effective amount of a pharmaceutical composition according to either claim 21 or 22.

36. A method for treating HIV infection in a mammal comprising the step of administering to said mammal a pharmaceutically effective amount of a pharmaceutical composition according to claim 23.

37. The method according to either claim 33 or 35, further comprising the step of concurrently or sequentially administering to the mammal one or more additional agents selected from the group consisting of other anti-viral agents and immunostimulators.

- 101 -

38. The method according to claim 37, wherein said other anti-viral agent or agents are protease inhibitors or reverse transcriptase inhibitors.

39. The method according to claim 38, wherein said protease inhibitor or inhibitors are HIV protease inhibitors.

40. The method according to claim 39, wherein said HIV protease inhibitor or inhibitors are selected from the group consisting of saquinavir (Ro 31-8959), MK 639, ABT 538 (A80538), AG 1343, XM 412, XM 450 and BMS 186,318.

41. The method according to claim 38, wherein said reverse transcriptase inhibitor or inhibitors are nucleoside analogs.

42. The method according to claim 41, wherein said nucleoside analog or analogs are selected from the group consisting of zidovudine (AZT), dideoxycytidine (ddC), didanosine (ddI), stavudine (d4T), 3TC, 935U83, 1592U89 and 524W91.

43. The method according to claim 38, wherein said reverse transcriptase inhibitor or inhibitors are non-nucleoside analogs.

44. The method according to claim 43, wherein said non-nucleoside reverse transcriptase inhibitor or inhibitors are delavirdine (U90) or nevirapine.

# INTERNATIONAL SEARCH REPORT

International Application No.

PCT/US 96/05475

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C07D307/20 C07D307/66 A61K31/34 A61K31/16 C07D417/12  
C07D405/12 C07D409/14 C07D413/12 C07D409/12 C07D413/14  
C07D407/12

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO,A,94 05639 (VERTEX PHARMA ;TUNG ROGER D (US); MURCKO MARK A (US); BHISETTI GOV) 17 March 1994 see examples with A containing THF in Table I see abstract; claims; table I see page 256 - page 262; example 197; tables VII,VIII ---	1-44
X	WO,A,95 06030 (SEARLE & CO ;MONSANTO CO (US); VAZQUEZ MICHAEL L (US); MUELLER RIC) 2 March 1995 see abstract; claims; examples 16A,16B,; tables 16A-16N,21 --- -/--	1-44

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

### \* Special categories of cited documents :

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

\*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

\*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

\*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

\*&\* document member of the same patent family

Date of the actual completion of the international search

9 August 1996

Date of mailing of the international search report

21.08.96

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentaan 2  
NL - 2280 HV Rijswijk  
Tel. (+ 31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+ 31-70) 340-3016

Authorized officer

Paisdor, B



# INTERNATIONAL SEARCH REPORT

Int. Jonal Application No  
PCT/US 96/05475

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT		
Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
P,X	<p>WO,A,95 33464 (SEARLE &amp; CO ;BRYANT MARTIN L (US); POTTS KAREN E (US); SMIDT MARY) 14 December 1995 see abstract; claims; example 11 -----</p>	1-44

**INTERNATIONAL SEARCH REPORT**

Information on patent family members

International Application No

PCT/US 96/05475

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO-A-9405639	17-03-94	AP-A- 390	02-08-95
		AU-B- 4852093	29-03-94
		BG-A- 99540	30-11-95
		CA-A- 2143208	17-03-94
		CN-A- 1087347	01-06-94
		CZ-A- 9500587	13-12-95
		EP-A- 0659181	28-06-95
		FI-A- 951059	18-04-95
		HU-A- 71892	28-02-96
		JP-T- 8501299	13-02-96
		LT-A,B 917	25-11-94
		NO-A- 950876	08-05-95
		PL-A- 307858	26-06-95
		SK-A- 29395	13-09-95
-----			
WO-A-9506030	02-03-95	AU-B- 7669794	21-03-95
		EP-A- 0715618	12-06-96
-----			
WO-A-9533464	14-12-95	AU-B- 2651095	04-01-96
-----			